



## **Biochemical Properties and Potential Application of Proteases from Alkalophilic *Bacillus lehensis* G1**

### **Authors:**

Noorul Aini Sulaiman\*, Nor Muhammad Mahadi and Nur Zazarina Ramly

\***Correspondence:** noorulaini@gmail.com

**DOI:** <https://doi.org/10.21315/tlsr2019.30.2.3>

### **Highlights**

- Extracellular proteases produced from Malaysia's *Bacillus* strain.
- Investigations on biochemical properties were performed on these extracellular proteases.
- Interesting digestion activities were observed when tested against x-ray film, feathers and stained cloth.

## **Biochemical Properties and Potential Application of Proteases from Alkalophilic *Bacillus lehensis* G1**

<sup>1,2</sup>Noorul Aini Sulaiman\*, <sup>2</sup>Nor Muhammad Mahadi and <sup>1</sup>Nur Zazarina Ramly

<sup>1</sup>Food Biotechnology, Faculty of Science and Technology, Islamic Science University of Malaysia (USIM), Bandar Baru Nilai 71800, Nilai, Negeri Sembilan, Malaysia

<sup>2</sup>Malaysia Genome Institute, Jalan Bangi, 43000 Kajang, Selangor, Malaysia

**Publication date:** 18 July 2019

**To cite this article:** Noorul Aini Sulaiman, Nor Muhammad Mahadi and Nur Zazarina Ramly. (2019). Biochemical properties and potential application of proteases from Alkalophilic *Bacillus lehensis* G1. *Tropical Life Sciences Research* 30(2): 25–37. <https://doi.org/10.21315/tlsr2019.30.2.3>

**To link to this article:** <https://doi.org/10.21315/tlsr2019.30.2.3>

**Abstrak:** Pencirian enzim ekstraselular protease daripada bakteria Alkalophilic *Bacillus lehensis* G1 dari Malaysia telah dikaji. Enzim protease yang dirembeskan diuji pada agar susu skim 2%. Keputusan menunjukkan protease ekstraselular mampu mengekalkan aktiviti sehingga suhu 60°C di dalam julat pH yang luas iaitu 3 hingga 11 dengan suhu optimum pada 40°C dan pH optimum pada 7.0. Aktiviti enzim juga diperhatikan akan meningkat dengan penambahan beberapa ion iaitu Mn<sup>2+</sup>, Fe<sup>2+</sup>, Cu<sup>2+</sup>, Mg<sup>2+</sup> dan Co<sup>2+</sup>. Manakala aktiviti protease didapati sedikit direncat dengan kehadiran ion Ca<sup>2+</sup>, K<sup>+</sup> dan Ni<sup>2+</sup> dengan baki aktiviti sebanyak 85%, 81% dan 75%. Protease ekstraselular juga didapati serasi dengan beberapa cecair detergen komersial dari Malaysia, yang menunjukkan protease ini boleh dimanfaatkan sebagai pembersih kotoran pada pakaian. Selain itu, potensi kegunaan protease yang dihasilkan oleh *B. lehensis* G1 ke atas penguraian gelatin dari filem X-ray yang telah digunakan juga telah dilakukan di dalam kajian ini.

**Kata kunci:** Alkalofilik, *Bacillus lehensis* G1, Protease, Pembersih Kotoran, Detergen, Filem X-ray, Penguraian Gelatin

**Abstract:** The biochemical properties of extracellular proteases enzymes from Malaysia Alkalophilic's bacteria, *Bacillus lehensis* G1, were investigated. The secreted enzymes were tested on 2% of skim milk agar. Results demonstrated that the enzyme could maintain the activity up to 60°C within an extensive range of pH from 3 to 11 with the optimal pH and temperature of 7.0 and 40°C, respectively. The proteases activity were also observed to be increased in the presence of several ions such as Mn<sup>2+</sup>, Fe<sup>2+</sup>, Cu<sup>2+</sup>, Mg<sup>2+</sup> and Co<sup>2+</sup>. Whilst, the enzyme activity was marginally inhibited with the addition of Ca<sup>2+</sup>, K<sup>+</sup> and Ni<sup>2+</sup> with the residual activity of 85%, 81%, and 75%, respectively. Furthermore, the extracellular proteases have shown to be compatible with several Malaysia commercial liquid detergents, which could be beneficial for stain removal. The potential application of proteases in gelatine decomposition from used X-ray films was also determined in this article.

**Keywords:** Alkalophilic, *Bacillus lehensis* G1, Proteases, Stain Removal, Detergent, X-ray Film, Gelatine Decomposition

---

\*Corresponding author: [noorulaini@gmail.com](mailto:noorulaini@gmail.com)

## INTRODUCTION

In Malaysia, the production of enzyme used to be considered almost non-existence, however, since Malaysian recognise the importance of enzymes, the development for enzyme industry becomes one of the national priorities in the area of biotechnology (Hii *et al.* 2012). Among the popular industrial enzymes is a protease. Protease is used as an additive in a variety of goods and processes such as in detergent, leather industry, photographic and biological-waste decomposition, medical treatment as well as the food industry (Bratanis *et al.* 2017; Baweja *et al.* 2016; Ray 2012).

Protease is an enzyme that hydrolysed the peptide bonds in amino acid residues of proteins or themselves and carried a vital role in various metabolic pathway such as in the process of immune system cascade amplification, vegetative growth, sporulation, activation of zymogen, as well as maintenance of the cellular protein pool (Esakkiraj *et al.* 2016; Li *et al.* 2013). Microorganisms generally produce a large array of protease, in the form of extracellular and/or intracellular (Jisha *et al.* 2013). Although proteases are continually produced worldwide, this enzyme is still insufficient to meet the growing demands of the industries. Thus, there has been a continuing exploration to find the new microbial sources of proteases with specific properties to fulfill the industrial enzyme requirements and to test their potential utilisation in various industry (Jayakumar *et al.* 2012; Lagzian & Asoodeh 2012; Anbu *et al.* 2015).

Novel extracellular proteases from *B. lehensis* G1 that was isolated from a soil sample in rubber estate, Johor, Malaysia (Sulaiman *et al.* 2017), has yet to be investigated. *Bacillus* is an aerobic or facultatively anaerobic growth, Gram-positive, endospore-forming bacteria with a huge diversity of physiological properties and is mostly studied for industrial applications (Cihan *et al.* 2012; Jayakumar *et al.* 2012). Whilst, *B. lehensis* G1 is known as alkali-tolerant and able to degrade various type of substrates including cellulose, starch, agar and hydrocarbons (Blanco *et al.* 2012). In this study, an effort has been made to describe the biochemical properties of *B. lehensis* G1's proteases followed with the potential use of the enzyme in gelatine decomposition and detergent.

## MATERIALS AND METHODS

### Media and Culture Conditions

Horikoshi agar (Blanco *et al.* 2012) was adjust to pH 10 (using sterile 1 M Na<sub>2</sub>CO<sub>3</sub>) was used to routinely grow the *B. lehensis* G1. The extracellular proteases from *B. lehensis* G1 were produced in Horikoshi medium, incubated at 37°C and 250 rpm for 24 h. The culture medium was centrifuged at 13,000 rpm for 10 min at 4°C, and the cell-free supernatant was then concentrated using a centrifugal filter unit (Amicon Ultra, Millipore). The cell-free supernatant was used as the crude enzyme was stored at -20°C until further used.

## Diffusion Plate Assay

Diffusion plate assay was performed by streaking *B. lehensis* G1 colonies on skim milk agar (2% Agar + 1% skim milk powder in 0.1 M Glycine-NaOH pH 10.0 buffer). The plate was incubated for 24 h and was observed for the development of halo-zone due to skim milk hydrolysis on the surface of the agar.

## Biochemical Properties of *B. lehensis* G1's proteases

A standard enzyme assay was used to measure the *B. lehensis* G1's proteases activity in pH 10.0 of Glycine-NaOH buffer according to Joshi and Satyanarayana (2013). The protein concentration was measured using the Bio-Rad protein assay kit based on the Bradford technique with bovine serum albumin (BSA) as standard. One unit of *B. lehensis* G1's proteases activity is defined as the total of protease required to release 1  $\mu$ mol of tyrosine per minute in the standard reactions.

The optimum pH of *B. lehensis* G1's proteases were examined in a different pH of buffers which includes Citrate (pH 3–5), Tris (pH 6–8), and Glycine-NaOH (pH 9–pH 11), respectively. Whilst, the effect of pH on the stability of *B. lehensis* G1's proteases activity were examined by pre-incubating the enzyme in the designated pH for 24 h. The residual activity was calculated using the standard enzyme assay.

The optimal temperature of *B. lehensis* G1's proteases were examined at different temperatures ranging from 25°C–90°C, and the enzyme activity was measured by standard assay procedure. Whilst, the temperature stability of the *B. lehensis* G1's proteases were carried out by pre-incubated the enzyme for an hour under the same temperature conditions.

## Effect of Metals and Surfactants

Effect of metal ions on *B. lehensis* G1's proteases activity was determined using  $\text{MnSO}_4$ , NaCl,  $\text{CuCl}_2$ ,  $\text{FeSO}_4$ ,  $\text{NiCl}_2$ ,  $\text{CoCl}_2$ ,  $\text{ZnCl}_2$ , KCl,  $\text{CaCl}_2$  and  $\text{MgSO}_4$  at 5 mM of concentrations. For the inhibitory effect on *B. lehensis* G1's proteases, divalent chelator and surfactants such as ethylenediaminetetraacetic acid EDTA (1 mM), sodium dodecyl sulphate (SDS), Triton-X-100, Tween 20, and Tween 80 were used (1% v/v). Crude proteases were pre-incubated with aforementioned metal ions and surfactants at 37°C for 1 h and then assayed for the residual activity using the standard assay.

## Protease Identification


Liquid Chromatography-tandem Mass Spectrometry (LC-MS/MS) was carried out at the Malaysia Genome Institute (National Institute of Biotechnology Malaysia). The peptide fragmentation results were given in FASTA sequence format was further subjected to align with *B. lehensis* G1's database ([http://www.mgi-nibm.my/bacillus\\_lehensis\\_g1/](http://www.mgi-nibm.my/bacillus_lehensis_g1/)).

## Potential Application of *B. lehensis* G1's Proteases


The extracellular proteases from *B. lehensis* G1 were tested against the removal of gelatine on used X-ray films, its compatibility with commercial detergents and washing capability on blood-stained cloth. The used X-ray films were cut into small pieces (1 cm x 1 cm) prior to incubating with crude proteases. The total of gelatine release into the solution was assayed at a 1 h time interval. For compatibility test; five commercial liquid detergents in Malaysia, such as Dynamo<sup>®</sup>, Breeze<sup>®</sup>, 5 CARE<sup>®</sup>, Tesco<sup>®</sup> and Tops<sup>®</sup> were boiled at 100°C for 10 min to deactivate their enzymes, and it was mixed with *B. lehensis* G1's proteases. The boiled detergents were initially assayed using standard protease assay to ensure that their enzyme was degraded. Then, the mixture of boiled detergents and *B. lehensis* G1' proteases were assayed for their residual activity and *B. lehensis* G1's proteases without detergents served as the control. Furthermore, cotton cloth was cut into pieces (4 cm x 4 cm) and stained with human blood. The cloth was allowed to air dry for 30 min. The removal of blood-stained was carried out by washing treatments using *B. lehensis* G1's proteases mixed with Tesco<sup>®</sup> liquid detergent at 40°C for 15 min using tap water. The result was visually measured. Stained cloth without added proteases served as the control.

## RESULTS AND DISCUSSIONS

An initial search of proteases produced by the *B. lehensis* G1 was tested using skim milk agar. The colonies of *B. lehensis* G1 showed substantial halo-zones which indicated the extracellular proteolytic activity on the skim milk substrate (Fig. 1). Then, the crude proteases were produced, followed by the examination of the enzyme properties. Activity assay of the extracellular proteases at different temperatures have shown that the optimal temperature of the enzyme was at 40°C. However, the enzymes still retained 65% of their activity at 50°C, as shown in Fig. 2(A). This finding showed that *B. lehensis* G1's proteases displayed a basal level of activity even at high temperature, which was between 70°C–80°C. Nevertheless, a further increase in the temperature caused a significant loss of the proteases activity. The thermos-stability of proteases was tested at different temperatures for an hour. Results showed that the enzymes were stable at room temperature and their optimal temperature with 100% activity remained (25°C–40°C). However, as shown in Fig. 2(B), the enzyme activity was decreased significantly from 20.1 to 4.95 U/mL when incubated at 60°C for 1 h. The temperature stability that is showed from *B. lehensis* G1's proteases is one of the important feature for an enzyme to be used in detergent industry (Pathak & Deshmukh 2012).



**Figure 1:** Skimmed milk agar plate showing secreted protease from *B. lehensis* G1 colonies; the zone of clearance around the colonies indicated proteolytic activity.



**Figure 2:** Optimal and stability tested for temperature and pH for extracellular proteases from *B. lehensis* G1. (A) optimum temperature, (B) thermo-stability, (C) pH optimum, and (D) pH stability. All the experiments were carried out in triplicate, standard error of  $8.088E-003$ ,  $p$ -value of 0.0042, and standard deviation of 19.73. The error bars in the experiments indicate standard deviation ( $n = 3$ ).


The optimum temperature of proteases from different *Bacillus* strains and species has been reported to be varied. Three different *B. cereus* strain NS-2, SIU1, and TKU022 have demonstrated diverse optimal temperature with 40°C, 45°C and 50°C, respectively (Singh *et al.* 2012; Liang *et al.* 2012; Bajaj *et al.* 2013). Several extracellular proteases showed a high optimal temperature at 60°C, such as *B. cereus*, *B. licheniformis* KBDL4 and *B. koreensis* BK-P21A (Saleem *et al.* 2012; Anbu 2013; Pathak & Deshmukh 2012). Whilst, extracellular protease

from hyperthermophilic *Bacillus* sp. MLA64 showed the highest protease activity at 95°C (Lagzian & Asoodeh 2012). Another study also showed that *B. circulans*, *B. clausii* and *B. licheniformis* have a high optimal temperature which is 85°C, 70°C and 67°C, respectively (Lagzian & Asoodeh 2012; Benkiar et al. 2013).

The effect of pH on the *B. lehensis* G1's proteases activity was tested at various pH using appropriate buffers. Results exhibited that the proteases reached the highest activity at neutral pH and decreased of activity towards either lower or higher pH (Fig. 2(C)). However, the enzymes were noticeably stable at different pH as the enzymes could retain nearly half of their activity at extremely low and high pH. The graph in Fig. 2(D) showed that the proteases were stable at pH 6–8 with slightly decreased of the activity at the value of 20%. The residual activity was recorded at 81%, 77% and 61% at pH 8, 9 and 10, respectively. Generally, serine protease is classified as an alkaliphilic enzyme in several *Bacillus* species. For example, the serine protease from *B. brevis* MWB-01 – pH 8.0, *B. koreensis* BK-P21A – pH 9.0, *Bacillus* sp. MLA64 – pH 9.5, *B. circulans* DZ100 – pH 12.5 and *B. lehensis* – pH 12.8 (Olajuyigbe & Falade 2014; Joshi & Satyanarayana 2013; Lagzian & Asoodeh 2012; Benkiar et al. 2013; Anbu 2013). Meanwhile, metalloprotease is known as a neutral protease. Wang et al. (2013) have shown metalloprotease from *B. amyloliquefaciens* SYB-001 are neutral protease with an optimal pH of 7.0. This is similar to metalloprotease from *B. stearothermophilus* and *B. thuringiensis* (Ou & Zhu 2012; Luo et al. 2013). This data is similar to our results of which metalloprotease behaved well at neutral pH. The neutral pH exhibited by *B. lehensis* G1 could be due to the abundant of metalloproteases as compared to the serine protease that is found to be secreted in the crude sample as showed by the mass spectrometry data. However, there is an exception for *B. cereus* TKU022 metalloprotease which prefers an alkali environment at pH of 10.0 (Liang et al. 2012).

According to Bajaj and Jamwal (2013), metal ions are essential for certain enzymes to carry out their full catalytic activity. Metal ions could affect the enzymes in several manners which includes (i) accepting or donating electrons to activate electrophiles or nucleophiles, (ii) acting as electrophiles, (iii) preventing unwanted side reactions, (iv) forming coordinate bonds between enzyme and substrate, (v) holding the reacting groups in the specific three-dimensional conformation and (vi) stabilizing the catalytically active site of the enzyme. The effect of metal ions on *B. lehensis* G1's proteases were tested with the final concentration of 5 mM and the total activity as shown in Fig. 3(A). Interestingly, ion  $Mn^{2+}$  increased the proteases activity at 2.5-fold higher (254%), followed by  $Cu^{2+}$ ,  $Fe^{2+}$  and  $Co^{2+}$  with 182%, 137% and 131%, respectively. Whilst, ion  $Ni^{2+}$ ,  $Zn^{2+}$ , and  $K^+$  caused a slight reduction in proteases activity. Meanwhile, proteases activity remained unchanged with the addition of ion  $Mg^{2+}$ . Furthermore, EDTA that was known as the metalloprotease inhibitor (Olajuyigbe & Falade 2014) was used in this study. The result has shown that *B. lehensis* G1 proteases have significantly loss their activity with the addition of 5 mM EDTA as shown in Fig. 3(A). This confirmed that there is an abundance of metalloproteases secreted by *B. lehensis* G1 as shown by the mass spec data. Surfactants such as Triton X-100, Tween 20, Tween 80,

and SDS was also tested in this study. These surfactants are used to test the stability of proteases activity. In this work, the enzymes were not entirely affected by these surfactants as showed in Fig. 3(B). The crude enzyme showed high stability in the presence of SDS, retaining 100% of its activity after incubation at 37°C for 1 h. Whilst, the residual activity of 80% and 78% retained by the enzyme with the addition of Tween 80 and Tween 20, respectively. However, incubation of enzyme with Triton X-100 significantly affected the enzyme with half of activity is inhibited. Similar result was showed by protease from *B. licheniformis* KBDL4 with high stability against an-ionic, non-ionic and oxidizing surfactant such as SDS and Tween 80. The stability of protease with the presence of surfactants is an important feature for detergent protease since the enzyme must be compatible and stable with all commonly used detergent compound such as SDS and Tween 80 that might be presence in the detergent formulation (Pathak & Deshmukh 2012).



**Figure 3:** Effect of (A) cation and (B) surfactant for extracellular proteases from *B. lehensis* G1. All the experiments were carried out in triplicate, standard error of 8.088E-003,  $p$ -value of 0.0431, and standard deviation of 21.90. The error bars in the experiments indicate standard deviation ( $n = 3$ ).


In mass spectrometry, results showed there was a total mixture of 168 proteins includes enzymes, chaperones, ribosomal and uncharacterised proteins in the extracellular protein of *B. lehensis* G1. Whilst, identification of protease revealed six metalloproteases and a single serine protease in the sample. The identified proteases in the secreted crude sample and its properties were summarised in Table 1. According to the MEROPS database (<https://www.ebi.ac.uk/merops/>), the *B. lehensis* G1's metalloproteases were subgroup into four

subtypes; M20, M23, M24, and M42 that are varied in their molecular structure, inhibitors, and conserved motif. However, in general, these metalloproteases are 'co-catalytic' with binding two metal ions; zinc or calcium. The metal ion is involved in the enzyme's catalytic activity, and the maintenance of the enzyme's structure rigidity (Ou & Zhu 2012).

**Table 1:** Mass spectrometry results for extracellular proteases of *B. lehensis* G1.

No.	BleG1	Proteases name	Size (kDa)	Calculated pI	Type of proteases	MEROPS
1	3821	Peptidase	49.8	5.03	Metalloprotease	M20
2	2767	Aminopeptidase	38.6	5.40	Metalloprotease	M42
3	2462	Peptidase	38.8	5.06	Metalloprotease	M24
4	2675	Aminopeptidase	38.2	5.39	Metalloprotease	M42
5	1563	Tripeptidase	39.8	5.62	Metalloprotease	M20
6	3825	Oligopeptidase	67.4	5.57	Serine protease	S9A
7	3033	Metalloprotease	51.9	4.72	Metalloprotease	M23


The second objective of this research is to find the potential applications of the proteases from *B. lehensis* G1. The application of proteases in silver recovery from X-ray film, its compatibility with Malaysia's commercial detergents as well as stain removal, were described in this paper. Various conventional methods have been applied for silver recovery from used photographic or X-ray film which includes burning the films, silver oxidation and using strong chemical solution for stripping the gelatine layer. However, these methods cause environmental pollution, hazardous, time-consuming and expensive. As an alternative step to extract the silver from the used X-ray film, the protease is used to break and release the silver by peeling the base coating from the surface of the X-ray film (Lakshmi & Hemalatha 2016). The proteolytic activity of the proteases on the gelatine decomposition on the used X-ray film was examined by measuring the amount of protein released into the solution at one-hour time intervals. The hydrolysis of the gelatine was initially slow on the first hour of incubation. However, the activity started to increase upon further incubation, as gelatine layer started to diminish from the surface of X-ray film. The proteases activity in the solution was recorded at 7.23 U/mL at 1 h incubation. After 4 h of incubation, the activity was recorded at 11.23 U/mL and remained stationary, indicating that the hydrolysis of the remained gelatine (Fig. 4(A)). The hydrolysed gelatine in the solution was verified by the changes of solution turbidity and the clearing of X-ray film (Fig. 4(B)). The gelatine layer was almost hydrolysed entirely from the X-ray film after 4 h, and this is showed by the clearing of the X-ray film (Fig. 4(C)). According to Bholay *et al.* (2012), *B. pumilus* p1's alkaline protease was found to be able to degrade gelatine from waste photographic films with 0.4013 g of silver weight and thus provide benefits in silver recovery application. This is an alternative step to extract silver from X-ray film which is eco-friendly, non-hazardous, time and cost effective.




**Figure 4:** Application of proteases on removal of gelatine layer from X-ray films. (A) Protease activity through clearing of gelatine on the surface of X-ray film tested in 7 h, (B) Changes of solution turbidity as gelatin release into solution, (i) control, without enzyme, (ii) test subject, and (C) Clearing of gelatine on the surface of X-ray film (i) control without enzyme, (ii) test subject after 2 h of incubation, (iii) test subject after 4 h of incubation. The error bars in the experiments indicate standard deviation ( $n = 3$ ).

The detergents compatibility study of *B. lehensis* G1 proteases have shown excellent stability after incubation with commercial detergents tested such as Dynamo®, Breeze®, 5CARE®, Tesco® and Tops®. This then may be applied as an additional component in the detergents formula. Results showed that proteases activities were retained for more than 85% of all the detergents tested, after being incubated at 40°C for 1 h, as shown in Fig. 5. The effectiveness of the *B. lehensis* G1' proteases in removing the stain from stained cloth with human blood was showed in Fig. 6. From five commercial detergents washing performance tested, only Tesco's detergent showed the lowest cleaning capability and therefore, *B. lehensis* G1' proteases were added to improve the washing performance. The blood-stained cloth that was washed with Tesco's detergent showed visible traces of blood stain retained on the cloth. However, after *B. lehensis* G1 proteases were added to the Tesco's detergent during the washing procedure, the blood stain was completely disappeared from the cloth. A relatively good wash performance was also demonstrated by the alkaline protease from *B. pumilus* MP27 and *B. subtilis* NCIM 2724 with the de-staining temperature at 50°C and 37°C, respectively (Baweja *et al.* 2016; Parpalliwar *et al.* 2015). These results

showed that the proteases produced from *B. lehensis* G1 could be utilised as a detergent additive since it shows good stability in commercial detergent tested, thermostable and as well as having good washing performance.



**Figure 5:** Compatibility test of extracellular proteases with commercial detergents in Malaysia. All the experiments were carried out in triplicate, standard error of 8.088E-003,  $p$ -value of 0.031, and standard deviation of 19.80. The error bars in the experiments indicate standard deviation ( $n = 3$ ).



**Figure 6:** Clearing of blood stain on the cloth using Tesco® detergent and *B. lehensis* G1's proteases. (A) control (without washing step), (B) stained cloth in 10 min washed (Tesco® detergent only), and (C) effective clearing of stained cloth in 10 min washed (Tesco® detergent with added *B. lehensis* G1's proteases).

## CONCLUSION

In this article, the characterisation and application of the extracellular proteases from *B. lehensis* G1 were investigated. The extracellular proteases have properties such as an optimum temperature of 40°C, optimum pH at 7.0, high thermostability and low sensitivity to pH. The neutral pH optimal, stability towards temperature,

pH, and surfactants could make this enzyme potentially useful for industrial applications such as silver recovery and detergent industry. An evaluation of its potential for use in the food industry will be the subject of future work.

## ACKNOWLEDGEMENTS

The authors are grateful to the Science Fund from the Malaysia Ministry of Science, Technology and Innovation (MOSTI) (Grant Number: USIM/SF-UKM/FST/30/40214) for financial support of this work.

## REFERENCES

- Anbu P. (2013). Characterization of solvent stable extracellular protease from *Bacillus koreensis* BK-P21A. *International Journal of Biological Macromolecules* 56: 162–168. <https://doi.org/10.1016/j.ijbiomac.2013.02.014>
- Anbu P, Gopinath S C B, Chaulagain B P and Lakshmi Priya T. (2015). Microbial enzymes and their applications in industries and medicine 2014. *BioMed Research International* 2015: 1–3. <https://doi.org/10.1155/2017/2195808>
- Bajaj B K and Jamwal G. (2013). Thermostable alkaline protease production from *Bacillus pumilus* D-6 by using agro-residues as substrates. *Advances in Enzyme Research* 1(2): 30–36. <https://doi.org/10.4236/aer.2013.12003>
- Bajaj B K, Sharma N and Singh S. (2013). Enhanced production of fibrinolytic protease from *Bacillus cereus* NS-2 using cottonseed cake as a nitrogen source. *Biocatalysis and Agricultural Biotechnology* 2(3): 204–209. <https://doi.org/10.1016/j.bcab.2013.04.003>
- Baweja M, Tiwari R, Singh P K, Nain L and Shukla P. (2016). An alkaline protease from *Bacillus pumilus* MP 27: Functional analysis of its binding model toward its applications as a detergent additive. *Frontiers in Microbiology* 7: 1–14. <https://doi.org/10.3389/fmicb.2016.01195>
- Benkiar A, Jaouadi Z, Badis A, Rebzani F, Touioui B, Rekik H, Naili B, Zohra F, Bejar S and Jaouadi B. (2013). International biodeterioration biodegradation biochemical and molecular characterization of a thermo- and detergent-stable alkaline serine keratinolytic protease from *Bacillus Circulans* Strain DZ100 for detergent formulations and feather-biodegradation. *International Biodeterioration Biodegradation* 83: 129–138. <https://doi.org/10.1016/j.ibiod.2013.05.014>
- Bholay A D, More S Y, Patil V B and Niranjana P. (2012). Bacterial extracellular alkaline proteases and its industrial applications. *International Research Journal of Biological Sciences* 1(7): 1–5.
- Blanco K C, de Lima C J B, Monti R and Jr J M. (2012). *Bacillus lehensis* – an alkali-tolerant bacterium isolated from cassava starch wastewater : Optimization of parameters for cyclodextrin glycosyltransferase production. *Annals of Microbiology* 62(1): 329–337. <https://doi.org/10.1007/s13213-011-0266-x>
- Bratanis E, Molina H, Naegeli A, Collin M and Lood R. (2017). BspK, a serine protease from the predatory bacterium *Bdellovibrio bacteriovorus* with utility for analysis of therapeutic antibodies. *Applied and Environmental Microbiology* 83(4): 1–16. <https://doi.org/10.1128/AEM.03037-16>

- Cihan A, Tekin N, Ozcan B and Cokmus C. (2012). The genetic diversity of genus *Bacillus* and the related genera revealed by 16S rRNA gene sequences and andra analyses isolated from geothermal regions of Turkey. *Brazilian Journal of Microbiology* 43(1): 309–324. <https://doi.org/10.1590/S1517-83822012000100037>
- Esakkiraj P, Meleppat B, Lakra A K, Ayyanna R and Arul V. (2016). Cloning, expression, characterization, and application of protease produced by *Bacillus cereus* PMW8. *RSC Advances* 6: 38611–38616. <https://doi.org/10.1039/C5RA27671C>
- Hii K L, Yeap S P and Mashitah M D. (2012). Cellulase production from palm oil mill effluent in Malaysia: Economical and technical perspectives. *Engineering in Life Sciences* 12(1): 7–28. <https://doi.org/10.1002/elsc.201000228>
- Jayakumar R, Jayashree S, Annapurna B and Seshadri S. (2012). Characterization of thermostable serine alkaline protease from an alkaliphilic strain *Bacillus pumilus* MCAS8 and its Applications. *Applied Biochemistry and Biotechnology* 168(7): 1849–1866. <https://doi.org/10.1007/s12010-012-9902-6>
- Jisha V, Smitha R, Pradeep S, Sreedevi S, Unni K, Sajith S, Priji P, Josh M S and Benjamin S. (2013). Versatility of microbial proteases. *Advances in Enzyme Research* 1(3): 39–51. <https://doi.org/10.4236/aer.2013.13005>
- Joshi S and Satyanarayana T. (2013). Characteristics and applications of a recombinant alkaline serine protease from a novel bacterium *Bacillus lehensis*. *Bioresource Technology* 131: 76–85. <https://doi.org/10.1016/j.biortech.2012.12.124>
- Lagzian M and Asoodeh A. (2012). An extremely thermotolerant, alkaliphilic subtilisin-like protease from hyperthermophilic *Bacillus* sp. MLA64. *International Journal of Biological Macromolecules* 51(5): 960–967. <https://doi.org/10.1016/j.ijbiomac.2012.08.009>
- Lakshmi B K M and Hemalatha K P J. (2016). Eco-friendly recovery of silver from used X-ray films by an alkaline protease of *Bacillus Cereus* strain S8. *Frontiers in Environmental Microbiology* 2(6): 45–48. <https://doi.org/10.11648/j.fem.20160206.14>
- Li Q, Yi L, Marek P and Iverson B L. (2013). Commercial proteases: Present and future. *FEBS Letters* 587(8): 1155–1163. <https://doi.org/10.1016/j.febslet.2012.12.019>
- Liang T W, Hsieh J L and Wang S L. (2012). Production and purification of a protease, a chitosanase, and chitin oligosaccharides by *Bacillus cereus* TKU022 fermentation. *Carbohydrate Research* 362: 38–46. <https://doi.org/10.1016/j.carres.2012.08.004>
- Luo X, Chen L, Huang Q, Zheng, J, Zhou W, Peng D, Ruan L and Sun M. (2013). *Bacillus thuringiensis* metalloproteinase Bmp1 functions as a nematocidal virulence factor. *Applied and Environmental Microbiology* 79(2): 460–468. <https://doi.org/10.1128/AEM.02551-12>
- Olajuyigbe F M and Falade A M. (2014). Purification and partial characterization of serine alkaline metalloprotease from *Bacillus brevis* MWB-01. *Bioresources and Bioprocessing* 1(8): 1–10. <https://doi.org/10.1186/s40643-014-0008-6>
- Ou J F and Zhu M J. (2012). An overview of current and novel approaches for microbial neutral protease improvement. *International Journal of Modern Biology and Medicine* 2(1): 1–31.
- Parpalliwar J P, Patil P S, Patil I D and Deshannavar U B. (2015). Extraction of silver from waste x-ray films using protease enzyme. *International Journal of Advanced Biotechnology and Research* 6(2): 220–226.
- Pathak A P and Deshmukh K B. (2012). Alkaline protease production, extraction, and characterization from alkaliphilic *Bacillus licheniformis* KBDL4: A Lonar soda lake isolate. *Indian Journal of Experimental Biology* 50(8): 569–576.
- Ray A. (2012). Protease enzyme: Potential industrial scope. *International Journal of Technology* 2(1): 1–5.

- Saleem M, Rehman A, Yasmin R and Munir B. (2012). Biochemical analysis and investigation on the prospective applications of alkaline protease from a *Bacillus cereus* strain. *Molecular Biology Reports* 39(6): 6399–6408. <https://doi.org/10.1007/s11033-011-1033-6>
- Singh S K, Singh S K, Tripathi V R and Garg S K. (2012). Purification, characterization and secondary structure elucidation of a detergent stable, halotolerant, thermoalkaline protease from *Bacillus cereus* SIU1. *Process Biochemistry* 47(10): 1479–1487. <https://doi.org/10.1016/J.PROCBIO.2012.05.021>
- Sulaiman N A, Mahadi N M and Ramly N Z. (2017). Identification of proteolytic genes from *Bacillus lehensis* G1. *Journal of Engineering and Science Research* 1(2): 14–20. <https://doi.org/10.26666/rmp.jesr.2017.2.3>
- Wang J, Xu A, Wan Y and Li Q. (2013). Purification and characterization of a new metallo-neutral protease for beer brewing from *Bacillus amyloliquefaciens* SYB-001. *Applied Biochemistry and Biotechnology* 170(8): 2021–2033.