

Relationship between *Ganoderma ergosterol* Concentration and Basal Stem Rot Disease Progress on *Elaeis guineensis*

Muniroh Md Saad, Nusaibah SA and Sariah Meon

Faculty of Agriculture, Universiti Putra Malaysia, Serdang, Selangor, Malaysia

*Corresponding author: nusaibah@upm.edu.my

Running title: Basal stem rot disease detection via ergosterol

Abstract. Basal stem rot (BSR) is a devastating disease to Malaysian oil palm. Current techniques employed for BSR disease detection on oil palm are laborious, time consuming, costly, and subjected to accuracy limitations. An ergosterol detection method was developed, whereby it correlated well with the degree of infection in oil palm. This current study was designed to study the relationship between *Ganoderma* biomass, ergosterol concentration, BSR disease progress and to validate the efficiency of microwave assisted extraction (MAE) method for extraction of ergosterol compound. In addition, testing on the sensitivity of thin layer chromatography (TLC) analysis for detection of ergosterol was also the aim of the present study. The optimized procedure involved extracting a small amount of *Ganoderma*-infected oil palm root tissues suspended in low volumes of solvent followed by irradiation in a conventional microwave oven at 70 °C and medium high power for 30 s, resulting in simultaneous extraction and saponification. Based on the results obtained, MAE method may be effective in extracting low to high yields of ergosterol from infected oil palm roots demonstrating disease scale 2, 3 and 4. Positive relationship was observed between ergosterol content and inoculation period starting day 3 in the inoculated oil palm seedlings and hour 6 in germinated seeds. TLC analysis demonstrated a good correlation with HPLC quantification. Therefore, a semi-quantitative TLC analysis may be applied for handling a large amount of samples during onset field survey.

Keywords : *Ganoderma*, basal stem rot, HPLC, microwave assisted extraction, TLC, oil palm

Introduction

Oil palm (*Elaeis guineensis*) is a monocotyledon in the family Arecaceae (formerly Palmae) within the subfamily Coccoideae (Corley and Tinker, 2003). It is a major crop in the tropical areas, particularly in the Southeast Asia. Palm oil is used worldwide for the production of food products, cosmetics, pharmaceuticals, biodiesel and in oleo chemical industry (Kalam and Masjuki, 2002; Corley and Tinker, 2003; Turner *et al.*, 2008). Oil palm industry contributes to the Malaysian economy by triggering the development of country's rural areas (Chin, 2008). In

Malaysia, cultivation of oil palm has increased year by year with 1.5 million ha in year 1985 to 5.84 million ha in 2018 (Malaysian Palm Oil Board, 2018).

Oil palm is subjected to numerous devastating diseases such as BSR, vascular wilt, spear rot-bud rot, sudden wither and red ring (Corley and Tinker, 2003). However, BSR is the major disease encountered by Malaysian palms, which is caused by *Ganoderma* spp. (Idris *et al.*, 2011). Paterson, (2019) reported that the disease was increasingly in inland peninsular Malaysia and Sabah, Malaysia, and in some cases at high levels, whereas before, it had not been detected. Several attempts have been made to control BSR using various control methods; however to date, none of the methods gave promising results in management of *Ganoderma boninense*, the major causal pathogen of BSR disease (Ariffin *et al.*, 2000; Sanderson *et al.*, 2000; Susanto *et al.*, 2005). The exertion of handling this disease is due to the infected palms not showing any external symptoms on mature palms until advanced stage. When it comes to this stage, the infected trees may not be able to respond to any treatment given (Bivi *et al.*, 2016). At present, the most common way used to detect BSR is based on the foliar symptoms and appearance of basidiomata at the base of infected stem. However, by the time visible symptoms appear, the palms were already at the final stages of infection and usually half of the basal tissues have been killed by the fungus (Idris, 2009). An early detection of BSR disease could prolong the economic life span of a palm (Lim *et al.*, 1990). Therefore, enzyme-linked immune sorbent assays-polyclonal antibody (Idris and Rafidah, 2008) as well as polymerase chain reaction (PCR) based techniques involving *Ganoderma*-specific primers (Bridge *et al.*, 2000; Utomo and Niepold, 2000; Yamoaka, *et al.*, 2000) have been proposed as early detection methods of the disease. However, these methods are complicated and time consuming for early detection of the disease in oil palm fields. Moreover, there are some limitations with PCR technique that requires to be addressed before applying for detection of *Ganoderma* (Paterson, 2007a; Paterson *et al.* 2008; Paterson and Lima, 2009). For instance, PCR could be subjected to inhibition (Paterson, 2007a) and ELISA-PAB (ELISA) suffers from cross reactivity (Idris and Rafidah, 2008). Hence, a feasible early detection method of the disease is required and crucial to prolong palm's life span via available curative methods.

Ergosterol is part of cell wall component of a fungus which is exclusively found in higher fungi and absence in other organisms (Madonna *et al.*, 2001; Mille-lindblom *et al.*, 2006). It is essential to fungus and its absence results in the death of the fungus (Marpugo *et al.*, 1964) thus indicating live fungal biomasses. The detection of ergosterol as fungal biomarker is measured to be the method of preference (Parsi and Gorecki, 2006). Ergosterol has been successfully used to indicate fungal biomass in soil (Grant and West, 1986; Frostegard and Baath, 1996), pathogenic fungi in roots and cereal grains (Bindler *et al.*, 1988; Seitz *et al.*, 1977), saprophytic fungi in decaying plant materials (Newell *et al.*, 1988), ectomycorrhizal fungi in roots and soil (Salmanowicz and Nylund, 1988; Wallander *et al.*, 1997), and recently in oil palm tissues (Mohd Aswad *et al.*, 2011; Toh Choon *et al.*, 2011; Chong *et al.*, 2014; Bivi *et al.*, 2016). Several attempts have been made to determine the relationship of ergosterol and fungal biomass under various conditions (Newell, 1992).

First data published on the use of ergosterol analysis as a diagnostic method to detect BSR supports the view that ergosterol exhibits the effectiveness for the detection of BSR in oil palm (Mohd Aswad *et al.*, 2011). Parkinson and Coleman (1991) reported that ergosterol assay was commonly considered the most promising tools for detection and quantification of fungal biomass. Parsi and Gorecki, (2006) also reported that the detection of ergosterol as fungal biomarker could be considered to be the method of choice.

Previous studies applied organic solvent based methods such as non-alkaline extraction (NAE) (Mohd Aswad *et al.* 2011), alkaline extraction (AE) (Zill *et al.* 1988), and ultrasonicator extraction (USE) (Yuan *et al.* 2007) methods for extraction of ergosterol. These organic solvent based methods (conventional) typically requires large samples size, large reagent volume, it is labor intensive and time consuming, additionally AE and USE were reported to yield low concentration of ergosterol compared to NAE method (Mohd Aswad *et al.*, 2011). Therefore, an efficient extraction method is required for the extraction of ergosterol. Young (1995) has developed MAE method for ergosterol extraction which requires a smaller sample size and reagent volume, it is more economical in terms of chemical used, and using convectional equipment (Domestic Microwave). Microwave assisted extraction is therefore more convenient than other methods in terms of time for sample preparation, cost and sample size. In addition, a larger sample size could be extracted at one time; moreover, the method is simple, rapid, and reliable for ergosterol detection on palms in the field environment during census carried out on disease survey.

Hence, the present study was undertaken to establish relationship between *Ganoderma* biomass, ergosterol concentration, and BSR disease progress in germinated seeds, artificially inoculated oil palm seedlings, and infected oil palm field tissues. Additionally, validation on the efficiency of MAE method for extraction of ergosterol and to test the sensitivity of TLC analysis for detection of ergosterol in artificially inoculated germinated seeds, artificially inoculated oil palm seedlings and infected oil palm field tissues was another objective.

Materials and Methods

Mycelial culture of *Ganoderma*

A pure culture of *G. boninense* was isolated from a basidiomata of an infected oil palm trunk in Felda Gua Musang, Malaysia using *Ganoderma* selective medium (GSM) (Ariffin and Idris, 1991).

Molecular identification

Molecular identification was conducted to confirm the *Ganoderma* culture. DNA was extracted using the modified CTAB method of Doyle and Doyle (1987). PCR Amplification was done as described by Utomo and Niepold (2000) with some modification on annealing temperature and amplification cycle. The PCR mixture containing 12.5 µl of Ampoun PCR master mix, 1 µl of both forward and reverse *Ganoderma* specific primers (Gan 1: 5' TTG ACT GGG TTG TAG

CTG 3' and Gan 2: 5' GCG TTA CAT CGC AAT ACA 3') (Utomo and Niepold, 2000) and 9.5 µl of nucleus free water were prepared in a 24 µl reaction volume. Then, 1 µl of DNA template was added to a final volume of 25 µl. The thermo cycler was programmed as follows: 5 min at 95°C, 35 cycles of 35 second at 94°C, 35 second s at 59.2°C, 40 second at 72°C, and 10 min at 72°C. The PCR products were analyzed by electrophoresis on a 1.5% agarose gel and stained with ethidium bromide to visualize the amplicates under UV light. The molecular identification from PCR product were sequenced using 1st BASE DNA sequencing services and aligned using BLAST network services against National Center for Biotechnology Information (NCBI

Quantification of ergosterol

Ergosterol extraction

Extraction of ergosterol from roots tissue were carried out using microwave assisted ectraction method based on the procedure by Muniroh *et al.*, (2014). 1.0 g of oil palm's root tissue was macerated in liquid nitrogen using a mortar and pestle into a powder, and transferred to a Pyrex test tube with a Teflon screw cap. Two mL of methanol (Merck, Chromatography grade) and 0.5 mL of 2M sodium hydroxide was added and the tube was tightly sealed. The test tubes were placed in a culture jar at the center of a conventional microwave (Sharp Jet Convectional Grill, model TTAG A437 with capacity 1.5 cu. ft) and subjected to microwave setting of 70°C, and medium high power with 30sec exposure time. The solutions were left to cool and were neutralized with concentrated hydrochloric acid. Finally, the solutions were extracted three times with 2 mL of pentane (Fisher chemicals, analytical reagent grade). The combine pentane extracts were then evaporated to dryness by using a Buchi rotary evaporator and then dissolved in 500 µl methanol for detection of ergosterol using TLC and quantified using HPLC with an ergosterol standard (Sigma, purity ≥ 95.0%).

Semi quantitative thin layer chromatography

Thin layer chromatography (TLC) was carried out to detect the presence of ergosterol from the extracted root tissues. TLC detection was undertaken based on Mohd Aswad *et al.*, (2011) in duplicate for all samples. The R_f value was calculated using this formula:

$$R_f = \frac{\text{Distance travelled by the product}}{\text{Total distance travelled by the solvent}}$$

High Performance Liquid Chromathography (HPLC)

An Agilent 1100 series HPLC equipped with a Diode Array Detector (G1315B), a pump (G1311A), and an auto sampler (G1313A) was used for quantification of ergosterol using an Ascentis express 2.7 µ C18 reverse-phase column (Supelco, USA). Operating conditions was based on Mohd Aswad *et al.*, (2011).

Germinated oil palm seeds

Germinated seeds (Dura x Pisifera) used were supplied by Sime Darby Research Center, Banting, Selangor. They were maintained in sterilized sand containing Hoagland solution for two weeks to allow rooting.

The experimental design for this experiment was complete randomized design (CRD) consists of two experimental treatments: non-inoculated with *G. boninense* (T1) and inoculated with *G. boninense* (T2). The germinated seeds were uprooted carefully and rinsed with distilled water. Germinated seeds for treatments (T2) were placed into 45-culture jar containing MAE slant culture of *G. boninense* (isolated from Gua Musang Felda) with three roots per culture jar. Non-inoculated germinated seeds (culture jar containing only MEA) were used as negative control. Random samplings of the experimental materials were done over a period of 6, 12, 24, 48, 72, 96, 120, 144 and 168 hours after inoculation with five culture jars per sampling time. All the roots were pooled and subjected to detection and quantification of ergosterol and further confirmed with PCR using modified method by Utomo and Niepold (2000). The roots sample from 0, 6, 24 and 48 hours were also subjected to scanning electron microscope (in-house method, Microscope Unit, Institute of Bioscience, UPM) to view the physiological changes after inoculation.

Oil palm seedling

The experiment was repeated using 6 month old oil palm seedling (Dura x Pisifera) supplied by Sime Darby Research Center, Banting, Selangor. The seedlings were maintained in polybags in the glasshouse until five to six leaf stages. The seedlings were watered daily and fertilized with NPK fertilizer (10 g per polybag) at monthly interval.

Eighty of 6 months-old oil palm seedlings were used for the infection study conducted in a glasshouse with two experimental treatments; non-inoculated with *G. boninense* colonized rubber wood block (RWB) (T1) and inoculated with *G. boninense* colonized RWB (T2). The seedlings were uprooted carefully and transplanted into polybags (size 12 cm x 15 cm) containing 3 kg soil mixture (3:2:1 v/v/v topsoil: peat: sand). Treatment (T2) was inoculated with a *G. boninense* mycelium colonized RWBs placed in contact with the roots (Sariah *et al.*, 1994). Non-inoculated seedlings were used as negative control. All oil palm seedlings were placed and arranged in a randomized complete block design (RCBD) under glasshouse conditions for 28 weeks. The seedlings were watered twice daily. Random destructive sampling of the seedlings was carried out on day 3, week 1, week 2, week 4, week 8, week 12, week 16, week 20, week 24, and week 28 with five replicates for each sampling. The root samples were used for detection and quantification of ergosterol.

A visual assessment of BSR infection was determined by examining the roots and foliar symptoms of the seedlings. The seedlings were also split longitudinally to observe root and bole decay and to visually assess the severity of the symptoms based on the proportion of number of

lesion (rotting) roots. The estimation was based on the scale modified from Breton *et al.*, (2005) (Table 1). Disease severity (DS) for internal and external symptoms of roots tissues and foliar symptoms was calculated based on formula derived from Liu *et al.* (1995) as follows:

$$\text{DS (internal)} = \frac{\sum \text{Number of seedlings in the rating} \times \text{rating number}}{\text{Total number of seedling assessed} \times \text{highest rating}} \times 100$$

$$\text{DS (External)} = \frac{\sum \text{Number of seedlings in the rating} \times \text{rating number}}{\text{Total number of seedlings assessed} \times \text{highest rating}} \times 100$$

Infected oil palm field tissues

Oil palm tissue samples were collected from high BSR disease incidence plot at Serting Felda Plantation Berhad, Negeri Sembilan. Mature palms aged 13 years old were randomly chosen based on the appearances of external symptoms of BSR disease and were categorized into scale 1, scale 2, scale 3, and scale 4 with 15 palms for each category (Table 2). Tissue samples (1.0 g) were weighed and grinded using liquid N₂ in a mortar and a pestle into fine powder. Samples were subjected to MAE extraction, TLC and HPLC analysis.

Statistical analysis

The data were analyzed using SAS Release 6 (SAS Institute Inc., 1990). Triplicate determinations of ergosterol concentrations from each sample were analyzed using ANOVA and means were compared by Least Significant Difference (LSD) ($P \leq 0.05$). Correlation analysis was performed using Microsoft Excel 2007.

Results

Identification of mycelial culture of *Ganoderma*

Nucleic acid of *Ganoderma* mycelium extracted using CTAB method was further identified using molecular identification. Polymerase chain reaction amplification with *Ganoderma* specific primer Gan1 and Gan2 were analyzed with 1.5% agarose followed by ethidium bromide staining showed visible band on the expected region at 150-200 bp (Fig. 1). Gene bank database

confirmed the samples were highly similar to *Ganoderma boninense* strain FA5017 with 99% similarity.

Germinated seeds

Ergosterol was detected in different concentrations in all inoculated germinated oil palm seeds and were absent in the un-inoculated germinated seeds (healthy samples). These concentrations apply for all the sampling periods determined from hour 6 to hour 168 after inoculation. Ergosterol was detected by visual evaluation image of TLC plates under UV-light in all inoculated seedlings (Sup fig. 1).

The R_f values of all samples were similar with that ergosterol standard spot with the value of 0.68. However, the ergosterol spot intensity under UV-light was faint and no detectable differences in inoculated seedlings at 6, 12, 24, 48, 72, and 96 hours after inoculation. However, the spot intensity increased gradually at hours 120, 144 and 168 after inoculation. High performance liquid chromatography analysis showed that ergosterol concentration increased with the increase of inoculation period (Fig. 2).

Ergosterol was detected as early as hour 6 after inoculation. Ergosterol concentration was significantly different from each sampling time from hour 6 to hour 168 after inoculation, however hour 120 and hour 144 did not show any significant differences after inoculation. The highest ergosterol concentration was 8.24 ug g^{-1} at hour 168 after inoculation with *G. boninense* culture, while the lowest ergosterol concentration was 0.96 ug g^{-1} on hour 6 after inoculation. A good correlation was observed between the inoculation period and ergosterol concentration ($R^2 = 0.97$) (Fig. 3).

Nucleic acid extracted using CTAB method from oil palm germinated seeds was further confirmed using molecular identification. Deoxyribonucleic acid amplification with *Ganoderma* specific primer Gan1 and Gan2 were analyzed with 1.5% agarose followed by ethidium bromide staining showed visible band for the inoculated germinated seeds on the expected region at 150-200 bp, while no visible band was observed for the un-inoculated germinated seed (Sup Fig. 2). Gene bank database confirmed the samples were highly similar to *G. boninense* strain FA5017 with 99% similarity.

These results were further confirmed by SEM. Whereby, SEM demonstrated there were no hyphae of *Ganoderma* sp. detected on the root (Fig. 4 A). In contrast, inoculated roots showed hyphae volumes increased with the increasing time of the inoculation period (Fig. 4 B, C, and D). The hyphae volume was very low at hour 6 after inoculation, however the hyphae started to colonise the roots at hour 24 after inoculation, and fully colonised the roots at hour 48 after inoculation.

Oil palm seedlings

Ergosterol was detected in different concentrations in all inoculated oil palm seedling roots from the samplings of day 3 to week 28. Ergosterol was also detected via images of TLC plates under UV-light in all inoculated seedlings (Sup Fig. 3). The R_f values of all samples were similar to that ergosterol standard spot with the value of 0.68cm. However, the ergosterol spot intensity under UV-light was faint and no detectable differences was found in inoculated seedlings at day 3, 7, 14, and week 4. Nonetheless, the spot intensity increased steadily from week 8 to week 28.

High performance liquid chromatography analysis showed ergosterol concentration increased parallel with the in internal and external disease severity from week 4 to week 28 (Fig. 5).

However, from day 3, 7 and 14, HPLC analysis quantified small amount of ergosterol from the samples, although there was no internal and external disease symptoms observed. The highest ergosterol concentration was 28.22 ug g^{-1} on week 28 with 100% external and internal disease severity where most of the palms were already dead while the lowest ergosterol concentration was 1.04 ug g^{-1} on day 3 with no observation of external and internal disease severity. A parallel correlation was observed between the internal disease severity with the ergosterol concentration ($R^2= 0.95$) and external disease severity with the ergosterol concentration ($R^2= 0.85$) (Sup fig. 4 and Sup fig. 5) respectively.

The oil palm seedling inoculated with *G. boninense* showed the presence of mycelium on the surface of the roots causing lesion and rotting to the roots, and the damages were observed increasing from day 3 to week 28 after inoculation (Sup Fig. 6). In addition, the infected palm showed a sign of stunted growth (Sup Fig. 7) and lesion of bole (Sup Fig. 8) when compared to the healthy seedlings. These symptoms can be observed clearly as the infection progresses from day 3 to week 28 after inoculation. From this study, internal disease severity can be detected a month after inoculation followed by external disease severity which can be observed two month after the inoculation.

Detection of ergosterol from infected field palm

Ergosterol was detected from all sample categorized in scale 2, scale 3, and scale 4 and absent from the scale 1 palms when analysed via visual evaluation of images with RP-18 Silica coated TLC plates (Merck) in UV-light based on the ergosterol standard spot. (Fig. 6). The R_f value of all detected samples were 0.68 (n-hexane: ethyl acetate) which is similar to R_f value of the ergosterol standard spot. The intensity of the spot increased in samples from scale 4, where the disease symptoms appeared as the appearance of foliar symptoms and presence of fruiting body at base of trunk.

In the HPLC analysis, ergosterol was identified by comparison of the retention time of ergosterol standard in all field palm samples. The ergosterol peak was well resolved and eluted at average of 7 min and the UV absorbance spectrum of ergosterol was clearly detected at 282 nm. The average concentration of ergosterol for Scale 2,3, and 4 were 6.31 , 9.27 , and $22.65 \text{ } \mu\text{g g}^{-1}$

respectively (Fig. 7). The ergosterol concentration varied from each infected palm with the concentration (scale 2 1-15) ranging from 3.59–11.03 $\mu\text{g g}^{-1}$, (scale 3 1-15) 5.95–14.4 $\mu\text{g g}^{-1}$, and (Scale 4 1-15) 17.11- 39.33 $\mu\text{g g}^{-1}$.

Discussion

Ganoderma boninense ergosterol concentration in artificially inoculated oil palm seedlings increased directly with the increase of inoculation period from day 3 to week 28. This finding was supported with the finding of Mohd Aswad et al. (2011) and Toh Choon et al. (2011). This present study also carried out an experiment on oil palm germinated seeds to study the relationship between inoculation time and ergosterol concentration and also to observe the earliest ergosterol can be detected prior inoculation with *Ganoderma*. From the result obtained, ergosterol was detected as early as six hours after inoculation which means that ergosterol can be detected once the fungus start to colonize the root. Thus, indicates that *G. boninense* mycelial mass colonizing the oil palm roots increases in abundant as a sign of disease progression following the period of inoculation. Xue et al. (2006) conducted a study on the correlation between ergosterol content of soybean fungal pathogens; *Diaporthe phaseolorum*, causal agent of Phomopsis seed decay, and *Cercospora kikuchii*, causal agent of leaf blight and purple seed stain and biomass of these pathogens on the host plant. The findings of the present study was also inline with Xue et al. (2006) whom reported that biomass was manipulated by the varying incubation period and resulted in the linearity correlation between fungal dry mass and ergosterol content.

A strong relationship was recorded between *G. boninense*'s ergosterol concentration and oil palm disease severity both internally and externally in artificially inoculated oil palm seedlings. The current findings were supported by a study conducted by Frey et al. (1992) which reported that concentration of ergosterol in infected roots differ significantly from control plants. In addition, Mohd Aswad et al. (2011) reported that ergosterol concentration detected from inoculated oil palm seedlings increased significantly with the increased degree of root infection. In the present study, the highest ergosterol concentration detected was 28.22 $\mu\text{g g}^{-1}$ on week 28 with 100% root infection, while the lowest ergosterol concentration was 1.04 $\mu\text{g g}^{-1}$ on day 3 after inoculation with no visible root infection. Detection of ergosterol as early as day 3 after inoculation indicated rapid disease establishment by *G. boninense* in the root tissues. In support to this finding, Nusaibah et al. (2016) reported that symptoms of *Ganoderma* infection on artificial infected oil palm seedling roots were visible via Scanning Electron Microscopy (SEM), Transmission Electron Microscopy (TEM) and plant defense response against *Ganoderma* attack via plant metabolites were also identified after 24 hours of inoculation period. In contrary, Mohd Aswad et al. (2011) managed to detect ergosterol via HPLC in *G. boninense*

inoculated oil palm roots after 3 weeks. This result was in contrast with the present study, and this could be due to the sample preparation before extraction. Whereby in the present study, roots were not washed or surface sterilized prior to ergosterol extraction, which enabled ergosterol detection on the colonized root surface tissues. Ergosterol was found in high concentrations as a fungal cell wall component (Gessner, 2005). In this experiment, the soil was sterilized prior to artificial inoculation steps to confirm that the soil used is free from other fungal contaminants. The results also demonstrated that external disease severity can be observed as early as week 8 after inoculation. Similarly, a study by Idris *et al.* (2006) reported that the foliar symptoms can be observed as early as two months after root inoculation in germinated seedlings.

For the field palm tissue samples, ergosterol was not detected in palms categorized in scale 1 which showed no visible symptoms. However, HPLC quantifies a small amount of ergosterol from scale 2 palms described as asymptomatic palms adjacent to the infected palms. Asymptomatic palms does not mean that the palm was free from disease. Mazliham *et al.* (2007) reported that visible symptoms of *Ganoderma* infection occurs at later stages of infection. Therefore from this finding, MAE method can be used as an early detection method of *Ganoderma* infection. HPLC quantified the highest ergosterol concentration in oil palm samples from scale 4 which exhibited the most severe symptoms. No significant differences were detected in ergosterol concentrations from scale 2 and 3 of infection. The average ergosterol concentration quantified by HPLC for scale 2, scale 3 were , 6.31 and 9.27 $\mu\text{g g}^{-1}$ respectively. However, palms with severe disease level in scale 4 resulted in highest ergosterol concentration with 22.65 $\mu\text{g g}^{-1}$ which was significantly different from palms from scale 1, 2 and scale 3 . Basal stem rot disease was reported as a white wood rotting process involving growth of the fungus within the oil palm tissues vialignin and cellulose biodegradation (Paterson, 2007b). Cellulose may be degraded readily by many fungi to gain energy, whereas lignin is a much more recalcitrant organic polymers that requires more energy for degradation purpose. *Ganodema boninense* was identified as a white rot fungus (Paterson, 2007b) which can fully degrade the lignin component with progression of the disease infection. By doing so, the biomass grew with a consequent increase in ergosterol, lignocellulotic enzyme and a weakening of the oil palm. Hence, detection of ergosterol could quantify the amount of growth which is related to the damage of oil palm.

In this study, no ergosterol was detected in non-inoculated and healthy palms which indicates that the ergosterol detected in the disease palm is from *G. boninense*. . The non-inoculated germinated seeds and seedlings grown under the same condition with inoculated samples showed no ergosterol detection when assessed using TLC and HPLC. This result indicates that healthy oil palm seedlings do not produce ergosterol as sterol compound. In addition, Zaiton *et al.* (2008) reported that most common endophytic microorganism found in healthy roots of symptomless palms were endophytic bacteria, and bacteria do not produce ergosterol.

Therefore, ergosterol could be a useful biochemical marker in detection of BSR disease in oil palm.

In the field palms, the sampling technique utilized played an important role to avoid contamination from other fungi. The oil palm trunk were drilled 0.5 – 1.0 m from the base of the palm to avoid contamination of microbes from the soil. Besides that, the driller used also was sterilize with 70% ethanol. To eliminate the fungus from the surface of oil palm trunk, the trunks were first drilled into 1-2 cm depth, and the driller was sterilized again before drilling further into the trunk. In this present study, extraction of the oil palm tissues were conducted within a week after the sampling period to avoid any contamination or decomposes of the samples. Fresh samples have to be used to obtain reproducible ergosterol data, and try not to expose the extracts to direct sunlight for prolonged period. Frey et al. (1992) reported that preliminary study conducted have shown that ergosterol decomposes when the root samples are dried or exposed to the ultraviolet light. These findings were similar to that reported by Newel et al. (1988) where ergosterol also degrades during freezing or lyophilization process.

Results obtained from SEM showed the hyphae colonization on the root samples inoculated with the *G. boninense*. As the fungal biomass increase, the hyphae colonization also increases with increased period of inoculation. Therefore, the ergosterol concentration increase with the increasing inoculation period of time. Rees et al. (2009) examined the *G. boninense* mode of infection using light microscopy and TEM and reported that root infection occurred consequence to firm attachment of *Ganoderma* hyphae to the root surface either localized to the initial point of contact or sometimes the fungus completely colonizes the root at the point of contact. Mille-Lindblom et al. (2004) reported the ergosterol method as a major advance in the estimation of fungal biomass. Larsen et al. (2004) also stated that ergosterol is the principal membrane sterol of most fungi and commonly used for estimating living fungal biomass.

In germinated seeds, the results were further confirmed using molecular detection. Polymerase chain reaction technique was employed to identify *Ganoderma* species (Moncalvo *et al.*, 1995; Idris *et al.*, 2003; Chong *et al.*, 2011). PCR product analysed resulted in the amplification fragment between 150-200 bp. Nevertheless, Utomo and Niepold (2000) conducted PCR using Gan1 and Gan2 on diseased oil palm roots samples and obtained amplified fragment size of 167 bp. Polymerase chain reaction amplification using *Ganoderma* specific primer Gan1 and Gan2 identified the pathogen as *G. boninense* with 99% similarity when BLASTn analysis performed on GenBank database. From the results of molecular identification, we could confirm the presence of *G. boninense* in the inoculated samples. Therefore, it was proven that the ergosterol detected in the oil palm root samples were from *G. boninense* fungal pathogen.

This present study also showed that the result of TLC were similar to the results obtained from quantification of ergosterol by using HPLC. For the germinated seeds and seedlings, the spot intensity of the TLC detection increase with the increase of inoculation period. HPLC also quantified the amount of ergosterol increased with the increasing inoculation period of time. In the field palm tissues, the spot intensity of the ergosterol detected in TLC showed the highest intensity for samples in scale 4. However, no significant difference in the spot intensity detected

in palms from scale 2 and scale 3 of infection. This result correlates with the results obtain with HPLC quantification where the highest ergosterol concentration produced were palms in scale 4 which was significantly different from palms in scale 2 and 3, while for palms in scale 2 and 3 showed no significant difference statistically.

Conclusion

Therefore, from this result, we can conclude that TLC analysis correlated well with the HPLC quantification. Thus, TLC analysis can be used for detection of ergosterol on the field palms as it is a more convenient and can be carried out on site besides suitable for large field survey during the census. Seitz et al. (1977) also suggested that preparative TLC and spectrophotometry could be used to estimate ergosterol if HPLC equipment is not available. Furthermore, according to Naewbanij et al. (1984) TLC can detect ergosterol as low as $1 \mu\text{g g}^{-1}$. In addition, ergosterol concentration also demonstrated a positive relationship between *Ganoderma* biomass and BSR development in artificially inoculated germinated seeds and seedlings. Moreover, ergosterol may be detected as early as 6 hours and 3 days after inoculation on germinated seeds and seedlings respectively using MAE method. In addition, the comparison between different extraction method was conducted and has been reported in Muniroh et al. (2014) and showed that MAE was the most efficient compared to the NAE and USE methods. Therefore, the use of MAE method in extracting ergosterol is suitable for the detection of BSR disease in field palms.

Acknowledgment

The Fundamental Research Grant Scheme (FRGS), administered through the Ministry of Higher Education, Malaysia, funded this project (Grant No: 5524175).

References

- Ariffin D., and Idris AS. (1991). A selective medium for the isolation of *Ganoderma* from diseased tissues. In: Yusof, B. *et al.* (eds). Proceedings of the 1991 PORIM International Palm Oil Conference-Progress, Prospects and Challenges Towards the 21st Century-Module 1, Agriculture. Palm Oil Research Institute of Malaysia, Bangi, Selangor, Malaysia 517-519
- Ariffin D., Idris A., and Singh G. (2000). Status of *Ganoderma* in oil palm. In: Flood, J. *et al.*, (eds), *Ganoderma* Diseases of Perennial Crops. CAB International Publishing, Wallingford, UK 249-666
- Benhamou N., Chet I. (1996). Parasitism of sclerotia of *Sclerotium rolfsii* by *Trichoderma harzianum*: ultrastructural and cytochemical aspects of the interaction. *Phytopathology* 86:405–416
- Bindler GN., Piadé JJ., and Schulthess D. (1988). Evaluation of selected steroids as chemical markers of past or presently occurring fungal infections on tobacco. *Beitrag Zur Tabakforschung International* 14:127–134

- Breton F., Hasan Y., Hariadi Lubis Z., and de Franqueville H. (2005). Characterization of parameters for the development of an early screening test for Basal Stem Rot tolerance in oil palm progenies. In: Proceedings of Agriculture, Biotechnology and Sustainability Conference. Technological Breakthroughs and Commercialization- The Way Forward, PIPOC 2005 MPOB International Palm Oil Congress, 25-29 September 2005, Kuala Lumpur, Malaysia
- Bridge PD., Ogrady EB., Pilotti CA., and Sanderson FR. (2000). Development of Molecular Diagnostics for the Detection of *Ganoderma* Isolates Pathogenic to Oil Palm. In: Flood, J. *et al.* (eds), *Ganoderma* Diseases of Perennial Crops. CAB International Publishing, Wallingford, UK 225-234
- Bivi, M. S., Paiko, A. S., Khairulmazmi, A., Akhtar, M. S., & Idris, A. S. (2016). Control of Basal Stem Rot Disease in Oil Palm by Supplementation of Calcium, Copper, and Salicylic Acid. *The plant pathology journal*, 32(5), 396–406
- Chin PFK. (2008). Malaysian efforts in developing responsible practises in the Palm Oil Industry, edited version of the keynote address by the Malaysian Minister of Plantation Industries and Commodities at the World Sustainable Palm Oil Conference, London, 15 September 2008. Published in the Global Oils & Fats Business Magazine .Vol. 5, Issue No. 4
- Chong KP., Lum MS., Foong CP., Wong CMVL., and Atong M. (2011). First identification of *Ganoderma boninense* isolated from Sabah based on PCR and sequence homology. *Afr. J. Biotechnol.* 10:14718-14723
- Corley RHV., and Tinker PB. (2003). The Oil Palm. 4th edition. *Oxford, UK: Blackwell Publishing*
- Frey B., Buser HR., and Schüepp H. (1992). Identification of ergosterol in vesicular-arbuscular mycorrhizae. *Biology Fertility Soils*, 13: 229-234
- Frostegard A., Baath E. (1996). The use of phospholipid fatty acid analysis to estimate bacterial and fungal biomass in soil. *Biology and Fertility of soils*, 22:59-65
- Gessner MO. (2005). Ergosterol as a measure of fungal biomass. In: Graça, M.A.S., Barlocher, F. and Gessner, M.O. (Eds.), *Methods to study litter decomposition: a practical guide*. 189-196, Dordrecht, The Netherlands. *Springer*
- Grant WD., and West AW. (1986). Measurement of ergosterol, diaminopimelic acid and glucosamine in soil: evaluations indicators of microbial biomass. *J. Microbiol Meth* 6:47-53
- Idris AS. (2009). Basal stem rot in Malaysia - Biology, Economic Importance, Epidemiology, Detection and Control. Paper presented at the International workshop on awareness, detection and control of oil palm devastating diseases, Kuala Lumpur, Malaysia. 6 November 2009
- Idris AS., and Rafidah AR. (2008). Polyclonal antibody for detection of *Ganoderma*. MPOB Information Series 430, MPOB, Malaysia

- Idris AS., Ismail S., Ariffin D., and Ahmad H. (2003). Control of Ganoderma-Infected palm-development of pressure injection and field applications. MPOB Information Series No. 131
- Idris AS., Kushairi A., Ariffin D., and Basri MW. (2006). Technique for inoculation of oil palm geminated seeds with *Ganoderma*. MPOB Information Series, MPOB TT No. 314, June 2006, Malaysian Palm Oil Board, Bangi, Selangor, Malaysia. 4
- Idris AS., Mior MHAZ., Maizatul SM., and Kushairi A. (2011). Survey on status of *Ganoderma* disease of oil palm. Proc. Of the PIPOC 2011 International Palm Oil Congress – Agriculture Conference. MPOB, Bangi 235-238
- Kalam MA., and Masjuki HH. (2002). Biodiesel from palm oil- an analysis of its properties and potential. Biomass Bioenerg 23: 471-479
- Khairuddin H. (1990). Basal stem rot of oil palm: incidence, etiology and control. Selangor, Malaysia: Master of Agriculture Science thesis, Universiti Pertanian Malaysia, Selangor, Malaysia
- Larsen T., Axelsen J., and Ravn HW. (2004). Journal of Chromatography A, 1026, 301-304
- Lim K., Chuah J., Ho C. (1993). Effects of soil heaping on Ganoderma infected oil palms. In: Proceedings of the PORIM International Palm Oil Congress. Update and Vision (Agriculture) 735–738
- Liu L., Kloepper JW., and Tuzun S. (1995). Induction of systemic resistance in cucumber against bacterial angular leaf sport by plant growth promoting rhizobacteria. *Journal of Phytopathology*, 85: 843-847
- Madonna AJ., Voorhees KJ., and Hadfield TL. (2001). Rapid detection of taxonomically important fatty acid methyl ester and steroid biomarkers using in situ thermal hydrolysis/methylation mass spectrometry (THM-MS): implication for bioaerosol detection. *J Anal and Appl Pyrol* 61: 65-89
- Malaysia Palm Oil Board, (2018). Oil Palm Planted Area By State As At December 2018. <http://bepi.mpob.gov.my/index.php/en/statistics/area/189-area-2018/857-oil-palm-planted-area-as-at-dec-2018.html> Microscopy Unit, Institute of Bioscience, Universiti Putra Malaysia, 43400 UPM, Serdang Selangor
- Mille-Lindblom C., Von Wachenfeldt E., and Tranvik L.J. (2004). Ergosterol as a measure of living fungal biomass: persistence in environmental samples after fungal death. *Journal of Microbiol Meth* 59: 253-262
- Mille-Lindblom, C., Fischer, H. and Tranvik L.J. (2006). Antagonism between bacteria and fungi: substrate competition and a possible tradeoff between fungal growth and tolerance towards bacteria. 113: 233-242
- Mohd Aswad AW., Sariah M., Paterson RRM., Zainal Abidin MA., Lima N. (2011). Ergosterol analysis of oil palm seedlings and plants infected with *Ganoderma* . *Crop Prot* 30: 1438-1442
- Moncalvo JM., Wang HH., and Hseu RS. (1995). Phylogenetic relationships in *Ganoderma* inferred from the internal transcribed spacers and 25S ribosomal DNA sequences. *Mycologia* 87: 223-238

- Morpurgo G., Serlupi-Crescenzi G., Tecce G., Valente F., and Venettacci D. (1964). Influence of ergosterol on the physiology and the ultra structure of *Saccharomyces cerevisiae*. *Nature*, 201: 897-899
- Muniroh MS., Sariah M., Zainal Abidin MA., Lima N., and Paterson RRM. (2014). Rapid detection of *Ganoderma*-infected oil palms by microwave ergosterol extraction with HPLC and TLC. *J. Microbiol. Methods* 100, 143–147
- Naewbanij M., Seib PA., and Burroughs R. (1984). Determination of Ergosterol using thin-layer chromatography and ultraviolet spectroscopy. *Cereal Chem* 61(5): 385-388
- Newell SY. (1992). Estimating fungal biomass and productivity in decomposing litter. in *The fungal community*, eds Carroll G. C., Wicklow D. T. (Marcel Dekker, Inc. New York, N.Y), 2: 521–561
- Newell SY., Arsuffi TL., and Fallon RD. (1988). Fundamental procedures for determining ergosterol content of decaying plant material by liquid chromatography. *Appl and Environ Microbi* 54:1876–1879
- Nusaibah SA., Siti Nor Akmar A., Idris AS., Sariah M., and Mohamad Pauzi Z. (2016). Involvement of metabolites in early defense mechanism of oil palm (*Elaeis guineensis* Jacq.) against *Ganoderma* disease. *Plant Physiol and Bioch*, 109:156-165
- Parkinson D., and Coleman DC. (1991). Microbial communities activity and biomass. *Agriculture, Ecosystems and environment* 34: 3-33
- Parsi Z., and Górecki T. (2006). Determination of ergosterol as an indicator of fungal biomass in various samples using non-discriminating flash pyrolysis. *J Chromatogr A*, 1130:145-150
- Paterson, R. R. M. (2019). *Ganoderma boninense* Disease of Oil Palm to Significantly Reduce Production After 2050 in Sumatra if Projected Climate Change Occurs. *Microorganisms*, 7(1), 24.
- Paterson RRM., and Lima N. (2009). Mutagens manufactured in fungal culture may affect DNA/RNA of producing fungi. *J Appl Microbiol* 106:1070-1080
- Paterson RRM. (2007a). Internal amplification controls have not been employed in diagnostic fungal PCR hence potential false negative results. *J Appl Microbiol* 102: 1369-1376
- Paterson RRM. (2007b). *Ganoderma* disease of oil palm - a white rot perspective necessary for integrated control. *Crop Prot* 26:1369–1376
- Paterson RRM., Sariah M., Lima N., Zainal Abidin MA., and Santos C. (2008). Mutagenic and inhibitory compounds produced by fungi affect detrimentally diagnosis and phylogenetic analyses. *Current Bioactive Compounds* 4: 245–257
- Rao AK. (1990). Basal stem rot (*Ganoderma*) in oil palm smallholdings. In: Ariffin D, Jalani S, editors. *Proceedings of the Ganoderma Workshop*, Palm Oil Research Institute of Malaysia. 113–131
- Rees RW., Flood J., Hasan Y., Potter U., and Cooper RM. (2009). Basal stem rot of oil palm (*Elaeis guineensis*); mode of root infection and lower stem invasion by *Ganoderma boninense*. *Plant Pathol* 58: 982-989

- Salmonawicz B., and Nylund JE. (1988). High performance liquid chromatography determination of ergosterol as a measure of ectomycorrhiza infection in Scots pine. *Eur J Forest Pathol* 18: 291-298
- Sanderson FR., Pilotti CA., and Bridge PD. (2000). Basidiospores: Their influence on our thinking regarding a control strategy for Basal Stem Rot of oil palm. In: Flood, J. *et al.* (eds), *Ganoderma* Diseases of Perennial Crops. CAB International Publishing, Wallingford, UK 113-121
- Sariah M., Hussin MZ., Miller RNG., and Holderness M. (1994). Pathogenicity of *Ganoderma boninense* tested by inoculation of oil palm seedlings. *Plant Pathol* 43: 507–10
- Seitz LM., Mohr HE., Burroughs R., Sauer DB. (1977). Ergosterol as an indicator of fungal invasion in grains. *Cereal Chemistry* 54:1207-1217
- Susanto, A. Sudharto, P.S. and Purba, R.Y. (2005). Enhancing biological control of basal stem rot disease (*Ganoderma boninense*) in oil palm plantations. *Mycopathologia*, **159**: 153-157.
- Toh Choon RL., Sariah M., Siti Mariam MN. (2012). Ergosterol from the soilborne fungus *Ganoderma boninense*. Short communication; *J Basic Microbiol* 52: 608-612
- Turner EC., Snaddon JL., Fayle TM., Foster WA. (2008). Oil Palm Research in Context: Identifying the Need for Biodiversity Assessment. *PLoS ONE* 3(2):1572
- Utomo C., and Niepold F. (2000). Development of diagnostic methods for detecting *Ganoderma* infected oil palms. *Journal of Phytopathology* 148: 507-514
- Wallander H., Massicotte HB., Nylund JE. (1997). Seasonal variation in protein, ergosterol and chitin in five morphotypes of *Pinus sylvestris* L. ectomycorrhizae in a mature swedish forest. *Soil Biology and Biochemistry* 29: 45–53
- Xue HQ., Upchurch RG., and Kwanyuen P. (2006). Ergosterol as a quantifiable biomass marker for *Diaporthe phaseolorum* and *Cercospora kikuchii*. *Plant Disease* 90:1395-1398
- Yamoaka M., Hayakawa S., Tsukamoto M., Kurane R., Idris AS., Mohd Haniff H., and Ariffin D. (2000). Diagnosis of basal stem rot of oil palm by foliar analysis and PCR-based detection of *Ganoderma* in oil palm. In the Proceedings of the 23rd Malaysian Society for Microbiology Symposium, Langkawi, Kedah 19-20 November 2000 4
- Young JC. (1995). Microwave-assisted extraction of the fungal metabolite ergosterol and total fatty acids. *J. Agric. Food Chem.* 43 (11):2904-2910
- Yuan JP., Hai Wang J., Liu X., Cong Kuang H., and Yan Zhao S. (2007). Simultaneous determination of free ergosterol and ergosteryl esters in *Cordyceps sinensis* by HPLC. *Food Chem* 105: 1755-1759
- Zaiton S., Sariah M., Zainal Abidin MA. (2008). Effect of endophytic bacteria on growth and suppression of *Ganoderma* infection in oil palm. *Int. J. Agric. Biol.*, 10: 127–132
- Zill G., Engelhardt G., and Wallnifer PR. (1988). Determination of ergosterol as a measure of fungal growth using Si 60 HPLC. *Z Lebensm Unters Forsch (European Food Research Technology)*, 187: 246-249

Table 1: Disease severity scale

Scale	Internal DS (Root lesion)	External DS (Foliar symptoms)
0	Healthy: no damage	Healthy
1	<10% rotting of roots; bole lesion	Yellowing of lower leaves and formation of rhizomorph at base of bole
2	10-20% rotting of roots; bole lesion	Necrosis of lower leaves and emergence of button-like sporophore at base of bole
3	20-50% rotting of roots; bole lesion	More than 50% necrosis of leaves and production of sporophore at base of bole
4	> 50% rotting of roots; bole lesion	Total necrosis and production of sporophore at base of bole

Table 2. Description of external BSR symptoms based on disease category level

Disease scale	Signs and symptoms
Scale 1	Palms apparently normal and free from disease
Scale 2	Asymptomatic neighboring palms with the infected palms
Scale 3	Palms with the presence of fruiting body at base of trunk
Scale 4	Appearance of foliar symptoms and presence of fruiting body at base of trunk



Figure 1: PCR amplification of 14 days old *Ganoderma* mycelial culture. M: Marker; L1-L4: amplified band of 14 days old *Ganoderma* mycelial culture.




Figure 2: Ergosterol concentration of the germinated seeds in un-inoculated and inoculated germinated seeds. No ergosterol was detected in un-inoculated germinated seeds. Bars represent SE (standard error) of triplicate determinations




Figure 3: Relationship of inoculation period and ergosterol concentration of oil palm germinated seeds




Figure 4: Comparison between inoculated and non-inoculated root (A) non-inoculated root, (B) 6 hours after inoculation, (C) 24 hours after inoculation, (D) 48 hours after inoculation with *Ganoderma* culture




Figure 5: Ergosterol concentration and disease severity percentage in inoculated seedlings. Bars represent SE (standard error) of triplicate determinations. (D= day; W= week; DS= disease severity)




Figure 6: TLC analysis of ergosterol from field palms based on different level of external BSR infection. Lane 1: Ergosterol standard, Lane2-4: Scale 1 palms, Lane 5-7: Scale 2 palms, Lane 8-10: Scale 3 palms, Lane 11-13: Scale 4 palms. (Scale 1= Palms apparently normal and free from disease ; Scale 2= Asymptomatic neighboring palms with the infected palms; Scale 3= Palms with the presence of fruiting body at base of trunk; Scale 4= appearance of foliar symptoms and presence of fruiting body at base of trunk).




Figure 7: Average ergosterol concentration ($\mu\text{g g}^{-1}$) compared to degree of BSR external symptoms (Scale 1= Palms apparently normal and free from disease; Scale 2= Asymptomatic neighboring palms with the infected palms; Scale 3= Palms with the presence of fruiting body at base of trunk; Scale 4= appearance of foliar symptoms and presence of fruiting body at base of trunk). Bars represent \pm SE (standard error) of triplicate determination