



Construction of Novel Yeast Strains from *Candida tropicalis* KBKTI 10.5.1 and *Saccharomyces cerevisiae* DBY1 to Improve the Performance of Ethanol Production Using Lignocellulosic Hydrolyzate

Authors:

Jamaluddin, Nisa Rachmania Mubarik, Edy Listanto and Eny Ida Riyanti*

***Correspondence:** enyriyanti@gmail.com, enyi001@brin.go.id

Submitted: 9 May 2022; **Accepted:** 19 September 2022; **Early view:** 11 November 2022

To cite this article: Jamaluddin, Nisa Rachmania Mubarik, Edy Listanto and Eny Ida Riyanti (in press). Construction of novel yeast strains from *Candida tropicalis* KBKTI 10.5.1 and *Saccharomyces cerevisiae* DBY1 to improve the performance of ethanol production using lignocellulosic hydrolyzate. *Tropical Life Sciences Research*.

Highlights

- Novel strains of Fcs1 and Fcs4 were obtained with different *xyl1* and *pdv5* genes from the *Candida tropicalis* KBKTI 10.5.1 and *Saccharomyces cerevisiae* DBY1 using genome shuffling method.
- The xylose-glucose consumption of Fcs1 and Fcs4 mutant on lignocellulosic hydrolyzate increased compared to their parents.
- The ethanol productivity of the Fcs1 mutant increased by 16.67% - 27.27%, while the Fcs4 mutant reached 22.22% - 33.33% compared to the parental KBKTI 10.5.1 and DBY1 when using acid hydrolyzate

Construction of Novel Yeast Strains from *Candida tropicalis* KBKTI 10.5.1 and *Saccharomyces cerevisiae* DBY1 to Improve the Performance of Ethanol Production Using Lignocellulosic Hydrolyzate

¹Jamaluddin, ²Eny Ida Riyanti*, ³Nisa Rachmania Mubarik and ²Edy Listanto

¹Study Program of Biotechnology, Graduate School of IPB University, Dramaga, Bogor 16680, Indonesia.

²National Research and Innovation Agency (BRIN), Jl. Tentara Pelajar No 3A, Bogor 16111, Indonesia

³Department of Biology, Faculty of Mathematics and Natural Science, IPB University, Dramaga, Bogor 16680, Indonesia.

*Corresponding author: enyriyanti@gmail.com

Running head: Novel Yeast Strains to Ethanol Production

Abstract. Increased consumption of xylose-glucose and yeast tolerance to lignocellulosic hydrolyzate are the keys to the success of second-generation bioethanol production. *Candida tropicalis* KBKTI 10.5.1 is a new isolated strain that has the ability to ferment xylose. In contrast to *Saccharomyces cerevisiae* DBY1 which only can produce ethanol from glucose fermentation. The research objective is the application of the genome shuffling method to increase the performance of ethanol production using lignocellulosic hydrolyzate. Mutants were selected on xylose and glucose substrates separately and using RAPD analysis. The ethanol production using lignocellulosic hydrolyzate by parents and mutants was evaluated using a batch fermentation system. Concentrations of ethanol, residual sugars, and by-products such as glycerol, lactate, and acetate were measured using HPLC machine equipped with Hi-plex H for Carbohydrate column and a Refraction Index Detector (RID) detector. Ethanol produced by Fcs1 and Fcs4 mutants on acid hydrolyzate increased by 26.58% and 24.17% from parent DBY1, by 14.94% and 21.84% from parent KBKTI 10.5.1. In contrast to the increase in ethanol production on alkaline hydrolyzate, Fcs1 and Fcs4 mutants only experienced an increase in ethanol production by 1.35% from the parent KBKTI 10.5.1. Ethanol productivity by Fcs1 and Fcs4 mutants on acid hydrolyzate reached 0.042 g/L/h and 0.044 g/L/h. The recombination of the genomes of different yeast species resulted in novel yeast strains that improved resistance performance and ethanol production on lignocellulosic hydrolysates.

Keywords: Biofuels, Fermentation, Genetic Engineering, Lignocellulosic, Yeast

INTRODUCTION

Renewable and environmentally friendly energy sources such as biofuels from biomass is essential strategies for substituting the use of fossil energy (Morais *et al.* 2018). The use of bioethanol has been proven to reduce carbon emissions compared to gasoline and diesel. The International Energy Agency (IEA) reports that the use of bioethanol is estimated to reduce carbon emissions by about 2.1 gigatons per year by 2050 if it is produced stably (Jin & Sutherland 2016). Bioethanol can be produced from abundant materials and can regenerate plants containing glucose, starch, and lignocellulose. Bioethanol production from lignocellulosic is more desirable than glucose and starch because socio-economically, sugar or starch leads to competition with food and feed needs (Meneses *et al.* 2017; Robak & Balcerek 2018).

Many factors affect the productivity of ethanol production using lignocellulosic, including the lack of ability of microorganisms to ferment xylose-glucose and low viable in lignocellulosic hydrolyzate media (Bušić *et al.* 2018; Robak & Balcerek 2018). The bioethanol production from xylose-glucose still uses a separate reactor since its inability or slower for co-fermenting both xylose and glucose (Cardona & Sa´nchez 2007). In addition, hydrolysis proceses of lignocellulosic produce several inhibitor compounds such as furfural, HMF, vanillin, acetic acid, and formic acid which can decrease yeast viability to ethanol production (Riyanti & Listanto 2017; Sjulander & Kikas 2020).

C. tropicalis KBKTI 10.5.1 1 can ferment xylose to produce ethanol, while *S. cerevisiae* DBY1 produce high ethanol concentration from glucose. Efforts to increase the ability to produce ethanol from xylose-glucose co-substrate continue to be carried out. Expression of xylose isomerase from *Burkholderia cenocepacia* to *S. cerevisiae* was performed (Vilela *et al.* 2013) however, the resulting strain still consumes xylose very slowly compared to glucose and produces relatively low ethanol productivity. Efforts to increase the consumption of xylose-glucose have also been carried out with an adaptive evolution approach on media containing xylose (Vilela *et al.* 2015). Research with this approach has produced a yeast strain that ferment xylose-glucose quickly and almost simultaneously. However, reports on genetic improvement of yeasts to increase ethanol fermentation of xylose-glucose co-substrates and their tolerance to lignocellulosic hydrolysates very limited. The complexity of the regulation system of co-fermenting xylose-glucose to produce ethanol and the response of yeast to inhibitor compounds in lignocellulosic hydrolysates makes it challenging to engineer its metabolism by inserting and/or knocking out several genes responsible for the regulation system, and time consuming. Therefore, undirected mutation technology through the genome

shuffling method has the potential to be developed to obtain superior yeast strains for lignocellulosic hydrolyzate fermentation.

Genome shuffling is a metabolic engineering method based on interprotoplast fusion. Protoplast fusion causes random recombination at the chromosomal level and can result in rapid multiplication of mutations so that the new strains formed have very diverse phenotypic properties (Biot-Pelletier & Martin 2014). Several studies have reported that genome shuffling can increase temperature and ethanol tolerance in *S. cerevisiae* 2013 (Orosco *et al.* 2013) as well as increase productivity and ethanol tolerance in *S. cerevisiae* and *Pichia stipites* (Jetti *et al.* 2018). Therefore, potential new strains could be obtained from *C. tropicalis* KBKTI 10.5.1 and *S. cerevisiae* DBY1 to increase xylose-glucose fermentation and have high tolerance to lignocellulosic hydrolysates.

MATERIALS AND METHODS

Yeast Strains

C. tropicalis KBKTI.10.5.1 and *S. cerevisiae* DBY1 were isolated from traditional Indonesian fermented foods that were carried out at the Molecular Biology Laboratory of the Indonesian Center for Agricultural Biotechnology and Genetic Resources Research (Indonesia). KBKTI.10.5.1 isolate was able to produce 0.8 g/L ethanol from 20 g/L xylose, while DBY1 could produce 21.98 g/L ethanol from 50 g/L glucose for 96 hours of incubation.

Culture Preparation and Media Preparation

Yeast isolates were cultured on YPD agar (1% yeast extract, 2% peptone, 2% D-glucose 2% agar) pH 5 for 24 hours. Seed culture for fermentation were prepared on liquid YPD media and incubated overnight at 30°C with 150 rpm agitation. Seed culture, as much as 10% (v/v) was used for fermentation. Lignocellulosic hydrolyzate of elephant grass (*Pennisetum purpureum* Schum) was obtained from 10 g of lignocellulose dissolved in 50 mL of acid solution (3% H₂SO₄) and alkaline solution (3% NaOH) for 24 hours followed by high temperature and pressure treatment using autoclave. The pH of the lignocellulosic hydrolyzate was adjusted to 5 using HCL and NaOH solutions (Riyanti *et al.* 2020).

Genome Shuffling (GS)

The construction of the novel strain was carried out by transferring DBY1 genomic DNA into KBKTI.10.5.1 cells using the genome shuffling method (Zhang & Geng 2012). The GS step was carried out with slight modifications to the pretreatment solution using 0.1 M CaCl₂. GS

was carried out through electroporation technique using the MicroPulser™ 165-2100 electroporation system. A total of 100 µl of GS results were spread into YPX media (1% yeast extract, 2% peptone, 2% D-xylose, 2% agar) pH 5 and incubated for 3 days at 30°C. The growing colonies were purified and mutants were selected.

Mutant Selection

Parent and mutants were inoculated into YPX and YPG agar (containing 1% yeast extract, 2% peptone, 2% D-xylose/D-glucose, 2% agar pH 5) and incubated overnight at 30°C, 150 rpm. Cultures with 1 mL (OD₆₀₀ = 0.5) were inoculated into 9 mL of YPX and YPG media and incubated for 24 hours at 30°C, 150 rpm. The ability of wild types and mutants to produce ethanol from xylose and glucose was evaluated by measuring the concentration of ethanol and the remaining fermented liquid substrate using HPLC (Riyanti *et al.* 2020). Mutant selection was also carried out through genome profile analysis using the Random Amplified Polymorphic DNA (RAPD) method with primers in Table 1.

Table 1: Random Amplified Polymorphic DNA (RAPD) Primers.

Primers	Sequences (5'- 3')
OPB-12	-CCTTGACGCA-
OPX-03	-TGGCGCAGTG-
OPX-06	-ACGCCAGAGG-
P-20	-AGGAGAACGG-

Evaluation of Cell Morphology and Genes Responsible for Ethanol Metabolism of Selected Strains

The selected strain cells were grown in YPD broth at 30°C for 48 hours. Cell cultures were prepared in a haemocytometer plate (depth 0-1 mm 1/400 mm²) and observed using a microscope. Several genes responsible for ethanol production and xylose utilization such as *xyl1*, *adh1*, *xks1*, *pdc1*, *pdc5*, *pdc6* were confirmed qualitatively using PCR method. The primers used are presented in Table 2.

Table 2 Primer pairs of genes that play a role in ethanol metabolism

Primers	Sequences (5' - 3')	Sources
xy11-F	ATGTCTACTACTCCTACTATTCCTAC	(Kim <i>et al.</i> 2019)
xy11-R	TTAAACAAAGATTGGAATGTTGTCCC	
adh1-F	AGACGCGCATAACCGCTAGA	(Smidt <i>et al.</i> 2012)
adh1-R	TAAGATGTGCGCATCTTGGGA	
xks1-F	CCAGTGATATCGAGGATGAGATTAGTAC	(Richard <i>et al.</i> 2000)
xks1-R	CCAGTGATATCTGTACTTGTGTCAGGGCAT	
pd1-F	ACACCATCTTGGCTTTGGTC	(Kahar <i>et al.</i> 2017)
pd1-R	CGAAAGCTGGGAATTGAGTC	
pd5-F	CACGTTGTTGGTGTTCATC	(Kahar <i>et al.</i> 2017)
pd5-R	TCAGTGATCATGGCAGTGGT	
pd6-F	GGAGATTGACCCCAACAAGA	(Kahar <i>et al.</i> 2017)

Ethanol Fermentation, Hydrolyzed Sugar Analysis, Analysis of Fermentation Results

Parent and mutants were grown in YPXG broth media (1% yeast extract, 2% peptone, 2% D-xylose, 5% D-glucose) and elephant grass (*Pennisetum purpureum* Schum) lignocellulosic hydrolyzate medium (pH 5). Fermentation was carried out for 72 hours on YPXG media and 24 hours on lignocellulosic hydrolyzate media at 30°C, 150 rpm. Observations were made every 24 hours on YPXG media and 4, 8, 24 hours on lignocellulosic hydrolyzate media. The content of xylose-glucose resulted from hydrolyzate lignocellulosic and its products such as ethanol, glycerol, lactate and acetate were analyzed using HPLC Agilent Technologies 1260 Infinity with a Hi-plex H for Carbohydrate column and a Refraction Index Detector (RID) detector (Riyanti *et al.* 2020).

Kinetic Parameters and Data Analysis

Cell biomass was obtained from the conversion of the OD₆₀₀ value (Myers *et al.* 2013). The ability of parental and mutant yeasts to consume xylose-glucose was calculated using the following formula:

$$\text{Percentage of xylose – glucose consumption} = \frac{S_0 - S}{S_0} \times 100\%$$

where,

S₀ = Initial xylose-glucose concentration

S = Xylose-glucose concentration after fermentation

The results of each concentration of ethanol and cell biomass obtained were then used to calculate the kinetic parameters of ethanol production. The kinetic parameters of ethanol production include ethanol yield (Y_{p/s}), biomass yield (Y_{x/s}) and ethanol productivity (Q_p) (Moremi *et al.* 2020).

$$Y_{p/s} = \frac{P}{\Delta S}$$

$$Y_{x/s} = \frac{X}{\Delta S}$$

$$Q_p = \frac{P}{\text{Fermentation Time}}$$

where,

P = Fermented products

X = Cell biomass

ΔS = Concentration of consumed substrate

The RAPD data obtained is based on the amplification band scoring with a classification of "1" if there is an amplified band and "0" if there is no amplified band. The data were then analyzed using the NTSYS-pc program (Numerical Taxonomy and Multivariate Analysis System, Version 2.02i). All ethanol production performance data were analyzed by simple mathematics using the excel program.

RESULTS

Selection of Potential Mutants for Ethanol Production from Xylose and Glucose

KBKTI 10.5.1 and DBY1 are isolates that have been identified using ITS1-ITS4 primers and have identified genomic characteristics using eight Random amplified polymorphic DNA (RAPD) primers (Jamaluddin et al. 2021). Random amplified polymorphic DNA is a DNA polymorphism produced by amplification of random DNA segments with a single primer sequence. The use of RAPD can be applied to distinguish between two individual organisms in one species (Suharsono et al. 2020). Thus, the approach to selecting potential fusants was to measure the fermentability of xylose and glucose into ethanol and by using RAPD technology.

The construction of new strains using genome shuffling by transferring genomic DNA of DBY1 into KBKTI 10.5.1 cell produced mutants which increased in xylose consumption and ethanol production (Table 3). In the screening process using glucose and xylose substrates, variable sugar uptake and ethanol production were observed from the GS mutants. Mutant Fcs2 experienced a decrease in xylose consumption rate compared to the parent of KBKTI 10.5.1. While mutants Fcs1, Fcs3 and Fcs5 shows an increase in xylose consumption rate and Fcs4 mutant show an increase in ethanol production using xylose and glucose substrate (Table 3). This phenomenon illustrates that there is a change in the genetic component which changes in metabolism. This is supported by the different genomic profiles between parents and mutants based on RAPD analysis (Figure 1).

Table 3 Test results for parental and mutant yeast ethanol production using xylose and glucose media

Substrates ^a	Ethanol Production Performance	Yeast Isolate ^b						
		Sc	Ct	Fcs1	Fcs2	Fcs3	Fcs4	Fcs5
Xylose	Sugar consumption (g/L)	0.04	1.94	2.22	1.86	2.52	1.91	2.14
	Ethanol products (g/L)	0.00	0.32	0.32	0.39	0.24	0.39	0.24
	Substrate efficiency (%)	0.19	9.70	11.1	9.31	12.6	9.53	10.6
Glucose	Sugar consumption (g/L)	19.8	19.5	19.6	19.5	19.6	19.5	19.6
	Ethanol products (g/L)	9	3	5	6	0	6	2
	Ethanol products (g/L)	7.34	6.94	6.94	7.02	7.02	7.02	7.02
	Substrate efficiency (%)	99.1	97.6	98.2	97.7	98.0	97.7	98.1
	Substrate efficiency (%)	7	7	3	9	2	9	0

Notes: ^aThe concentration of substrates is 20 g/L fermented for 24 hours, ^b(Sc) *S. cerevisiae* DBY1, (Ct) *C. tropicalis* KBKTI 10.5.1, (Fcs) mutans strain




Figure 1. RAPD electrophorogram of parental and mutant yeast genomes on 1% agarose. (M) 1 kb Ladder DNA, (Sc) *S. cerevisiae* DBY1, (Ct) *C. tropicalis* KBKTI 10.5.1, (Fcs) mutant strains.

The RAPD profile in Fig. 1 illustrates that GS causes the transfer or substitution of nucleotide bases from DNA DBY1 to genomic DNA of KBKTI 10.5.1. The existence of insertion or substitution of nucleotide bases causes the RAPD primers used to attach or not during the PCR process so that polymorphisms are formed. The results of the RAPD analysis showed that there was random recombination between the genome of DBY1 and KBKTI 10.5.1. The RAPD profile of parental and mutant polymorphisms using the OPX-03 primer showed that the mutants similar to the KBKTI 10.5.1, but there was an additional DNA band at 2500 bp as shown in the DBY1 profile. RAPD profiles with P-20 primer were found different between parent and mutans as band around 150 bp was do not detected in Fcs1, also band around 250 pb was not detected in in Fcs 5. The difference in RAPD amplicon profiles as shown in Fig. 1 indicates the presence of genetic diversity in the constructed mutants. The resulting mutants were clustered in one cluster with the parent KBKTI 10.5.1. Fcs1 mutants were far apart to form separate sub-clusters at a genetic distance of 0.70. In addition, the Fcs4 and Fcs5 mutants grouped together to form a separate sub-cluster from the parent of KBKTI 10.5.1

(Fig. 2). The existence of genetic diversity produced can be an opportunity to produce superior strains. Fcs1 and Fcs4 mutants were then selected for further observation. Selection of Fcs1 and Fcs4 mutants based on the best ethanol production and xylose consumption compared to their parents.




Figure 2. Cluster dendrogram of parental and mutant yeast strains constructed with SAHN-UPGMA. (Sc) *S. cerevisiae* DBY1, (Ct) *C. tropicalis* KBKTI 10.5.1, (Fcs) mutant strains.

Genomic DNA of *C. tropicalis* KBKTI 10.5.1 has also been tried to be transferred into *S. cerevisiae* DBY1 cells, but the yeast strain obtained was still unable to produce ethanol from xylose (unpublished data). The failure of the construction of the new yeast strain for ethanol production from xylose using of strategy the KBKTI 10.5.1 genomic DNA transfer into DBY1 cells was thought to be because *S. cerevisiae* DBY1 does not have a xylose reductase gene sequence, a gene that plays a role in catalyzing xylose to enter ethanol metabolism (Mouro *et al.* 2020; Olofsson *et al.* 2011). Thus, the DNA sequence of the KBKTI 10.5.1 xylose reductase gene was not successfully inserted into DBY1 DNA because there was no homology between the two sequences.

Evaluation of Cell Morphology and Genes Responsible for Ethanol Metabolism of Selected Mutant

The novel yeast strains that were constructed had morphological characters of cells which were thought to be inherited from the two parents. The two mutant cells were spherical in shape more closely resembled parental KBKTI 10.5.1 compared to the slightly oval shape of DBY1 cells. However, the mutant cells did not form pseudohyphae like the parent of KBKTI 10.5.1 which had been incubated at 30°C for 48 hours simultaneously with the two parents (Fig. 3).




Figure 3. Morphology of parental and mutant yeast cells after 48 hours of incubation on YPD media. (a1, b1, c1 and d1) budding, (b2) pseudohyphae.

In addition to cell morphology observations, several genes that play a role in ethanol metabolism were also evaluated qualitatively. The Fig. 4 shows that KBKTI 10.5.1 and both mutants have the same amplicon xylose reductase profile, which was around 1000 bp. However, the results of the sequence alignment analysis showed that there were four nucleotide base differences (Fig. 5). This indicated that the transfer of DBY1 DNA into KBKTI 10.5.1 cells caused a change in the nucleotide base arrangement in the KBKTI 10.5.1 genome. The results of the analysis of sequence homology in the *BLAST-n* program showed that the xylose reductase KBKTI 10.5.1 was identical to the xylose reductase *Candida*

tropicalis isolate GRA1 (99.59%) while the Fcs1 mutant changed to 99.28% against the xylose reductase isolate GRA1 gene. *Candida tropicalis* isolate GRA1 is a comparison isolate, which has the gene sequences encoding xylose reductase available in the genebank (MF143598.1) Isolate GRA1 is able to reduce xylose to xylitol, which will enter the ethanol fermentation pathway, because it has a xylose reductase gene sequence (Ariyan and Uthandi 2019). The difference in the percentage similarity of the xylose reductase sequences of KBKTI 10.5.1 and Fcs1 isolates to GRA1 can be one of the benchmarks for the success of the genome shuffling that has been carried out. Changes in the arrangement of four nucleotide bases can change the amino acid composition of the mutant, this is indicated by the results of *BLAST-p* xylose reductase KBKTI 10.5.1 which has a homology percentage of 96.96% and Fcs1 xylose reductase reaches 96.27% against D-Xylose reductase *Candida tropicalis* (QED90344 .1).




Figure 4. Electrophorogram of genes that play a role in ethanol metabolism. (M) 1 kb Ladder DNA, (1) *S. cerevisiae* DBY1, (2) *C. tropicalis* KBKTI 10-5.1, (3) Fcs1, (4) Fcs4




Figure 5. The results of the Alignment analysis of the xylose reductase (*xyl1*) sequences of *C. tropicalis* KBKTI 10-5.1 and Fcs1 using the BioEdit program

The gene encoding pyruvate decarboxylase were also different. The results of the study in Fig. 4 show that there are several bands generated from the *pdc5* and *pdc6* PCR primers. This shows that the primer used is not yet specific. However, it can be seen that with *pdc5* and *pdc6* primers, Fcs1 and Fcs4 mutants are different from their parents. Differences in the amlicone profile of the gene encoding pyruvate decarboxylase may support differences in the production of by-products (lactate, acetate, and glycerol), as shown in Fig. 6.




Figure 6. The bioethanol production performance of parental and mutant yeast in YPXG media

Performance of Wild Type and Mutant Yeast for Ethanol Production on YPXG Media and Elephant Grass (*Pennisetum purpureum* Schum) Lignocellulosic Hydrolyzate

Yeast strains capable of fermenting xylose-glucose to produce high ethanol concentrations are one of the keys to the success of ethanol production using lignocellulosic (Zabed *et al.* 2016). Our results showed that the KBKTI 10.5.1 strain as well as the two constructed mutants had the ability to produce approximately 2% ethanol from xylose-glucose fermentation for 72 hours (Fig. 6). However, the ability to use xylose-glucose between parents and mutants was different. The DBY1, KBKTI 10.5.1 and the Fcs4 mutant were not able to use xylose-glucose simultaneously, in contrast to the Fcs1 mutant which was able to use xylose when glucose

was still available in the medium. In addition, the consumption of xylose by the Fcs1 mutant was slightly higher than that of the two parents, but there was no increase in ethanol production. This phenomenon is thought to occur because the consumed xylose was converted to lactate, this can be seen from the increase in lactate production as the xylose concentration in the media decreases (Fig. 6). However, Fcs1 and Fcs4 mutants are potential to be used as second generation bioethanol production agents with further research on process engineering to minimize the formation of side products such as lactate, acetate and glycerol.

The results of the study as shown in Fig. 6 can be seen that the use of xylose at a mixed concentration of 20-50 g/L xylose-glucose has not been able to significantly increase ethanol production. The research report showed that *Pichia kudriavzevii* and *C. tropicalis* experienced an increase of ethanol production at a 1:1 xylose-glucose mixture (Nweze *et al.* 2019). Therefore, it is necessary to optimize the concentration of the xylose-glucose mixture to increase ethanol production. In addition to substrate concentration, the ability to produce ethanol using lignocellulosic is also influenced by the presence of inhibitors produced in the hydrolysis process (Faizal *et al.* 2020; Chaudhary *et al.* 2021). Therefore, this study also reports the performance of ethanol by parental and mutant strains in lignocellulosic hydrolyzate media presented as in Figs 7 and 8.




Figure 7. The bioethanol production performance of parental and mutant yeast in elephant grass lignocellulosic acid hydrolyzate media




Figure 8. The bioethanol production performance of parental and mutant yeast in elephant grass lignocellulosic alkaline hydrolyzate media

Elephant grass (*Pennisetum purpureum* Schum) is one of the potential raw materials for bioethanol production (Vidal *et al.* 2017). Elephant grass biomass contains 60.20% cellulose, 23.80% hemicellulose (Minmunin *et al.* 2015), which can be hydrolyzed into xylose and glucose. However, the concentration of sugar (xylose-glucose) obtained from the hydrolysis of acids and bases in this study was very low, reaching only 4.18 g/L and 1.77 g/L (Figures 7 and 8). This can indicate that the hydrolysis of elephant grass biomass using acids and bases combined with pressure has not been maximized to increase the concentration of glucose and xylose. Sugar concentration below 10 g/L in bioethanol fermentation is not recommended because it can produce very little ethanol (Faizal *et al.*, 2020). However, the results of our study need to be reported because through the production of bioethanol using lignocellulosic hydrolysates, the information on the ability to consume xylose-glucose and the growth ability of yeast strains that we obtained on the hydrolyzate can be known.

The performance of xylose-glucose consumption between parents and mutants on both acid and alkaline hydrolyzate media showed that the two mutants were more similar to the parents of KBKT1 10.5.1. (Figs. 7 and 8). The ability to consume glucose by DBY1 on acid hydrolyzate media seemed to slow down, where at 4 hours glucose was still available compared to the other three strains that had been used up. This is one of the effects of the

concentration of ethanol produced being lower than the parent KBKTI 10.5.1 and the two mutants (Fig. 7). The results in Figure 7 also show that at the concentration of the xylose-glucose mixture of about 3:1, the parent of KBKTI 10.5.1 and both mutants were able to use xylose-glucose simultaneously. This is indicated by the significant use of xylose at 4 hours (Fig.7). In addition, the ability to grow mutants on acid hydrolyzate media was better when compared to the two parents, this was indicated by the high OD₆₀₀ value and a slight increase in ethanol production (Fig. 7).

Productivity of Parent and Mutant Yeast on Lignocellulosic Hydrolyzate Media

The ability of bioethanol production by the parents and the resulting new strains are different. This difference can also be seen from the kinetic parameters as shown in Table 4. The ethanol productivity of the parents on YPXG media with a composition of 20-50 g/L xylose was better than that of the mutant, but on acid hydrolyzate the ethanol productivity of the new strain was better. This can indicate that genomic recombination between the two parents resulted in improved productivity of resistance to lignocellulosic hydrolysates. The conversion of xylose-glucose to ethanol by strains Fcs1 and Fcs4 on acid hydrolyzate media was also quite efficient. This can be seen from the product yield which reached 0.37 g/g. The xylose-glucose conversion efficiency of strains Fcs1 and Fcs4 was almost equivalent to the conversion of hexose with the addition of a lignocellulosic inhibitor by *Saccharomyces cerevisiae* isolate ISO12, which was 0.38 g/g, in which the ISO12 strain was claimed to be one of the superior candidates for the production of second generation bioethanol (Wallace-Salinas & Gorwa-Grauslund 2013; Riyanti *et al.* 2020).

Table 4: Parameters of parental and mutant yeast kinetics on different media.

Isolate	Medium ^a	Fermentation Time (h)	Percentage of Xylose-Glucose Consumption (%)	Ethanol (g/L)	Yp/s (g/g)	Yx/s (g/g)	Qp (g/L/h)
Sc	YPXG	72	74.90	18.66	0.36	0.04	0.259
	HA	24	30.37	0.79	0.67	0.35	0.033
	HB	24	77.05	0.78	0.61	0.18	0.033
Ct	YPXG	72	76.79	18.08	0.33	0.03	0.251
	HA	24	69.60	0.87	0.31	0.23	0.036
	HB	24	100	0.74	0.45	0.25	0.031
Fcs1	YPXG	72	78.30	17.91	0.33	0.03	0.249
	HA	24	69.60	1.00	0.37	0.29	0.042
	HB	24	100	0.75	0.47	0.20	0.031
Fcs4	YPXG	72	77.63	17.94	0.33	0.03	0.249
	HA	24	71.76	1.06	0.37	0.29	0.044
	HB	24	100	0.75	0.46	0.22	0.031

^aThe concentration of xylose-glucose in YPXG media is 20-50 g/L, acid hydrolyzate (HA) media is 3.08-1.11 g/L and alkaline hydrolyzate (HB) is 0.47-1.30 g/L.

DISCUSSION

Genetic mutation is one of the phenomena that can cause the diversity of organisms. The use of the concept of mutation has developed rapidly, such as to assemble a superior organism (Galhardo *et al.* 2007; Loewe & Hill 2010). This study uses a random mutation approach in two different species with the help of electric shocks. The use of electric shocks aims to enlarge the pores of the cell membrane so as to increase the permeability of the membrane so that DNA molecules can enter the cell (Nesin *et al.* 2011). The electric shock will cause an unused break in the second molecule of the parent yeast genomic DNA. This disconnection provides an opportunity for mutations to occur when there is a homologous wet tide and an incorrect reunification. Mutations that occur can reach more than one gene so that the mutants obtained have diversity at the genetic level (Figures 1 and 2).

The constructed novel strain has a different physiology from the two wild types. Among the different physiological properties are the ability to produce ethanol from xylose, glucose and cause differences in cell morphology such as the formation of pseudohyphae. Pseudohyphae are formed from shoot cells, such as blastospores, which multiply, but the daughter cells do not separate from the parent cell and continue to elongate to resemble hyphae, so that there is a septum between the blastospore and the growing part of the cell, and in this section there is a narrowed part (Veses & Gow 2009). The formation of pseudohyphae is closely related to the physiological properties of cells which are influenced by several factors such as nitrogen deficiency (Lackey *et al.* 2013; González *et al.* 2018), CO₂ exposure (Sasani *et al.* 2016), high phosphate and growth period (Veses & Gow 2009), growing temperature conditions (Nadeem *et al.* 2013). The results of cell morphology observations as shown in Figure 3 can illustrate that the physiological roles of parent and mutant in responding to environmental conditions (nutrients in YPD media, temperature and incubation time) were different. The difference in phenotype of an organism is part of the influence of its genetic expression (Young *et al.* 2019). Therefore, the transfer of DBY1 DNA to KBKTI 10.5.1 cells was suggested to cause the insertion of genetic material into the KBKTI 10.5.1 genome, thereby causing changes in some genetic expressions.

The novel strain that was constructed experienced an increase in the ability to consume xylose and produce ethanol. Yeast strains capable of fermenting xylose cannot use xylose until the glucose concentration is completely depleted. This can be explained by several factors, including because glucose can suppress the expression of genes responsible for xylose metabolism through the transcription factor Mig1. Cells that are in high glucose concentrations cause Mig1 to move from the cytoplasm to the nucleus and bind to gene promoters that can cause repression of xylose catabolism genes. After the glucose concentration in the medium decreases, the transcription factor Mig1 is transported back to

the cytoplasm so that the repression is released (Rolland *et al.* 2002). This is one of the factors that slows down ethanol production in the presence of xylose in the media. High glucose concentration in the xylose-glucose mixture is one of strategy to increase ethanol production. This can be seen from the ability of the parent and mutant for 24 hours to produce ethanol around 6.94-7.34 g/L from the fermentation of 20 g/L glucose (Table 3), compared to a mixture of 20-50 g/L xylose-glucose already can produce approximately 2% ethanol or about 17-18 g/L (Fig. 6). Moreover, the study results need to be reported because the constructed mutants have an increased ability to consume xylose (Table 3) and can even be used in together with glucose (Figs. 6, 7 and 8).

The use of xylose-glucose by KBKTI 10-5.1 and the two mutants on elephant grass lignocellulosic hydrolyzate media showed that both were used simultaneously but were not able to maximally increase ethanol production (Figures 7 and 8). This is thought to be caused by the very low concentration of total sugars, so that the sugars consumed is mainly used up for cell growth. In addition, there is no inhibitors separation process was done prior the fermentation process, therefore complex inhibitors produces during acid and alkaline hydrolysis may adding the stress condition during fermentation process.

Yeast cells are the main engine in fermentation, having the potential to be exposed to various types of fermentation stresses such as osmotic stress, temperature stress, ethanol stress and the presence of toxic compounds during the fermentation process (Burphan *et al.* 2018; Mukherjee *et al.* 2014). The resistance of yeast cells in responding to fermentation stress is an important factor that must be possessed by a fermentation agent (Riyanti & Listanto 2020). There is a direct relationship between fermentation efficiency and yeast resistance to stress which refers to the ability of yeast lines to adapt efficiently to unfavorable growing conditions (Saini *et al.* 2017). Lignocellulosic hydrolyzate can produce several mixtures of inhibitor compounds that can inhibit ethanol production, cell growth, and enzyme biochemical activity (Palmqvist & Hahn-Hagerdal 2000). Therefore, OD₆₀₀ nm in Figures 7, 8, showed that the newly obtained lines experienced faster growth in the lignocellulosic hydrolyzate than the parent. This can represent the ability of tolerance and good productivity to the presence of toxic compounds lignocellulosic hydrolyzate.

Yeast cells that grow undergo a mechanism of nutrient transport and assimilation accompanied by the integration of nutrients into cellular components, so that there is an increase in biomass and cell division. The main purpose of growing yeast cells is to multiply cells rather than produce ethanol (Walker & Stewart 2016). However, during fermentation, ethanol production and yeast growth are closely related processes. Therefore, the ability to transport xylose-glucose hydrolyzate lignocellulosic is one of the factors that affect the productivity of ethanol production. The genomic recombination of KBKTI 10.5.1 and DBY1 succeeded in improving the genes responsible for xylose-glucose transport to cells, this was

indicated by an increase in xylose-glucose consumption in Fcs1 and Fcs4 mutants by 3.64% - 4.54% from DBY1 and 1.09% -1.97% from KBKTI 10.5.1 on YPXG media. An increase in xylose-glucose consumption also occurred in both acidic and basic hydrolyzate media (Table 4).

CONCLUSION

Novel strains of Fcs1 and Fcs4 were obtained with different *xy11* and *pd5* genes from the *Candida tropicalis* KBKTI 10.5.1 and *Saccharomyces cerevisiae* DBY1. The performance of xylose-glucose consumption and ethanol productivity of Fcs1 and Fcs4 mutant on lignocellulosic hydrolyzate increased compared to their parents. The ethanol productivity of the Fcs1 mutant increased by 16.67% - 27.27%, while the Fcs4 mutant reached 22.22% - 33.33% of the parental KBKTI 10.5.1 and DBY1 on acid hydrolyzate.

ACKNOWLEDGEMENTS

We would like to thank to Indonesian Centre for Biotechnology and Genetic resources Research and Development, Indonesian Agency for Agricultural Research and Development for funding this research. This research has been funded by register number DIPA:4584.SDA.503.052.A.

REFERENCES

- Ariyan M, and Uthandi S. (2019). Xylitol production by xylose reductase over producing recombinant *Escherichia coli* M15. *Madras Agricultural Journal* 106: 205–209. doi:10.29321/MAJ.2019.000247
- Biot-Pelletier D and Martin V J J. (2014). Evolutionary engineering by genome shuffling. *Applied Microbiology and Biotechnology* 98(9): 3877–3887.
- Burphan T, Tatip S, Limcharoensuk T, Kangboonruang K, Boonchird C and Auesukaree C. (2018). Enhancement of ethanol production in very high gravity fermentation by reducing fermentation-induced oxidative stress in *Saccharomyces cerevisiae*. *Scientific Reports* 8(13609): 1–11. <https://doi.org/10.1038/s41598-018-31558-4>
- Bušić A, Marđetko N, Kundas S, Morzak G, Belskaya H, Šantek M I, Komes D, Novak S and Šantek B. (2018). Bioethanol production from renewable raw materials and its

- separation and purification: A review. *Food Technology & Biotechnology* 56(3): 289–311.
- Cardona C A and Sa´nchez O J. (2007). Fuel ethanol production: process design trends and integration opportunities. *Bioresource Technology* 98(12): 2415–2457. <https://doi.org/10.1016/j.biortech.2007.01.002>
- Chaudhary A, Hussain Z, Aihetasham A, El-Sharnouby M, Rehman R A, Khan M A U, Zahra S *et al.* (2021). Pomegranate peels waste hydrolyzate optimization by response surface methodology for bioethanol production. *Saudi Journal of Biological Sciences* 28(9): 4867–4875.
- Faizal A, Sembada A A and Priharto N. (2020). Production of bioethanol from four species of duckweeds (*Landoltia punctata*, *Lemna aequinoctialis*, *Spirodela polyrrhiza*, and *Wolffia arrhiza*) through optimization of saccharification process and fermentation with *Saccharomyces cerevisiae*. *Saudi Journal of Biological Sciences* 28(1): 294–301.
- Galhardo R S, Hastings P J and Rosenberg S M. (2007). Mutation as a stress response and the regulation of evolvability. *Critical Reviews in Biochemistry and Molecular Biology* 42(5): 399–435.
- González B, Vázquez J, Cullen P J, Mas A, Beltran G and Torija M. (2018). Aromatic amino acid-derived compounds induce morphological changes and modulate the cell growth of wine yeast species. *Frontiers in Microbiology* 9(670): 1–16. <https://doi.org/10.3389/fmicb.2018.00670>
- Jamaluddin, Mubarik N R, Listanto E, and Riyanti E I. (2021). Isolation and Molecular Identification of Yeasts that Capable to Fermenting Xylose for Second Generation Bioethanol Production. *Proceesings: National Seminar of the National Commission Genetic Resources*. <http://repository.pertanian.go.id/handle/123456789/14483>
- Jetti K D, GNS R R, Garlapati D and Nammi S K. (2018). Improved ethanol productivity and ethanol tolerance through genome shuffling of *Saccharomyces cerevisiae* and *Pichia stipites*. *International Microbiology* 22(2): 247–254.
- Jin E and Sutherland J W. (2016). A proposed integrated sustainability model for a bioenergy system. *Procedia CIRP* 48(2016): 358–363. <https://doi.org/10.1016/j.procir.2016.03.159>
- Kahar P, Riyanti E I, Otsuka H, Matsumoto H, Kihira C, Ogino C and Kondo A. (2017). Challenges of non-flocculating *Saccharomyces cerevisiae* haploid strain against inhibitory chemical complex for ethanol production. *Bioresource Technology* 245(2017): 1436–1446. <https://doi.org/10.1016/j.biortech.2017.06.009>
- Kim S, Lee J and Sung B H. (2019). Isolation and characterization of the stress-tolerant *Candida tropicalis* YHJ1 and evaluation of its xylose reductase for xylitol production from acid pre-treatment wastewater. *Frontiers in Bioengineering and Biotechnology* 7(138): 1–12. <https://doi.org/10.3389/fbioe.2019.00138>

- Lackey E, Vipulanandan G, Childers D S and Kadosh D. (2013). Comparative evolution of morphological regulatory functions in *Candida* species. *Eukaryotic Cell* 12(10): 1356–1368.
- Loewe L and Hill W G. (2010). The population genetics of mutations: good, bad and indifferent. *Philosophical Transactions of the Royal Society B: Biological Sciences* 365(1544): 1153–116.
- Meneses L R, Raud M, Orupold K and Kikas T. (2017). Second-generation bioethanol production: A review of strategies for waste valorization. *Agronomy Research* 15(3): 830–847.
- Minmunin J, Limpitipanich P and Promwungkwa A. (2015). Delignification of elephant grass for production of cellulosic intermediate. *Energy Procedia* 79(2015): 220–22534. <https://doi.org/10.1016/j.egypro.2015.11.468>
- Morais R F, Morais C S B, de Morais L F and Almeida J C C. (2018). Energy balance of elephant grass biomass for power generation by direct biomass combustion. *African Journal of Biotechnology* 17(13): 405–410.
- Moremi M E, Rensburg E L J V and Grange D C L. (2020). The improvement of bioethanol production by pentose-fermenting yeasts isolated from herbal preparations, the gut of dung beetles, and marula wine. *International Journal of Microbiology* 2020: 1–13. <https://doi.org/10.1155/2020/5670936>
- Mouro A, Santos A A D, Agnolo D D, Gubert G F, Bon E P S, Rosa C A, Fonseca C and Stambuk B U. (2020). Combining xylose reductase from *Spathaspora arborariae* with xylitol dehydrogenase from *Spathaspora passalidarum* to promote xylose consumption and fermentation into xylitol by *Saccharomyces cerevisiae*. *Fermentation* 6(3): 72. <https://doi.org/10.3390/fermentation6030072>
- Mukherjee V, Steensels J, Lievens B, de Voorde I V, Verplaetse A, Aerts G, Willems K A, Thevelein J M, Verstrepen K J and Ruyters S. (2014). Phenotypic evaluation of natural and industrial *Saccharomyces* yeasts for different traits desirable in industrial bioethanol production. *Applied Microbiology and Biotechnology* 98(22): 9483–9498.
- Myers J A, Curtis B S and Curtis W R. (2013). Improving accuracy of cell and chromophore concentration measurements using optical density. *BMC Biophysics* 6(4): 1–15. <https://doi.org/10.1186/2046-1682-6-4>
- Nadeem S G, Shafiq A, Hakim A T, Anjum Y and Kazm S U. (2013). Effect of growth media, pH and temperature on yeast to hyphal transition in *Candida albicans*. *Open Journal of Medical Microbiology* 3(3): 185–192.
- Nesin O M, Pakhomova O N, Xiao S and Pakhomov A G. (2011). Manipulation of cell volume and membrane pore comparison following single cell permeabilization with 60- and 600-ns electric pulses. *Biochimica et Biophysica Acta Biomembranes* 1808(3): 792–801.

- Nweze J E, Ndubuisi I, Murata Y, Omae H and Ogbonna J C. (2019). Isolation and evaluation of xylose-fermenting thermotolerant yeasts for bioethanol production. *Biofuels* 12(8): 961–970. <https://doi.org/10.1080/17597269.2018.1564480>
- Olofsson K, Runquist D, Hahn-Hägerdal B and Lidén G. (2011). A mutated xylose reductase increases bioethanol production more than a glucose/xylose facilitator in simultaneous fermentation and co-fermentation of wheat straw. *AMB Express* 1(4): 1–8. <https://doi.org/10.1186/2191-0855-1-4>
- Orosco F L, Estrada S M, Simbahan J F, Alcantara V A and Pajares I G. (2017). Genome shuffling for improved thermotolerance, ethanol tolerance and ethanol production of *Saccharomyces cerevisiae* 2013. *Philippine Science Letters* 10(1): 22–28.
- Palmqvist E and Hahn-Hägerdal B. (2000). Fermentation of lignocellulosic hydrolysates. II: inhibitors and mechanisms of inhibition. *Bioresource Technology* 74(1): 25–33.
- Richard P, Toivari M H and Penttilä M. (2000). The role of xylulokinase in *Saccharomyces cerevisiae* xylulose catabolism. *FEMS Microbiology Letters* 190(1): 39-43.
- Riyanti E I and Listanto E. (2017). Inhibition of the growth of tolerant yeast *Saccharomyces cerevisiae* strain I136 by a mixture of synthetic inhibitors. *Indonesian Journal of Agricultural Science* 18(1): 17–24.
- Riyanti E I and Listanto E. (2020). Understanding yeast tolerance as cell factory for bioethanol production from lignocellulosic biomass. The Second International Conference on Genetic Resources and Biotechnology. Bogor, Indonesia, 24–25 May 2021, 060006-1–060006-7.
- Riyanti E I, Yuniawati R, Sanjaya R A, Samudra I M, Listanto E, Lestari E G and Mastur. (2020). Potential use of *Saccharomyces cerevisiae* TKPK 10.5.1 isolated from several sources for ethanol using various sugar sources. The 6th International Conference on Biological Science ICBS 2019. Yogyakarta, Indonesia, 10–11 October 2019, 060004-1–060004-9.
- Robak K and Balcerak M. (2018). Review of second generation bioethanol production from residual biomass. *Food Technology and Biotechnology* 56(2): 174–187.
- Rolland F, Winderickx J and Thevelein J M. (2002). Glucose-sensing and –signaling mechanisms in khamir. *FEMS Yeast Research* 2(2): 183–201.
- Saini P, Beniwal A and Vij S. (2017). Comparative analysis of oxidative stress during aging of *Kluyveromyces marxianus* in synthetic and whey media. *Applied Biochemistry and Biotechnology* 183(1): 348–361.
- Sasani E, Khodavaisy S, Agha K A S, Darabian S, Rezaie S. (2016). Pseudohyphae formation in *Candida glabrata* due to CO₂ exposure. *Current Medical Mycology* 2(4): 49–52.

- Sjulander N and Kikas T. (2020). Origin, impact and control of lignocellulosic inhibitors in bioethanol production—A Review. *Energies* 13(4751): 1–20. <https://doi.org/10.3390/en13184751>
- Smidt O, Preez J C and Albertyn J. (2012). Molecular and physiological aspects of alcohol dehydrogenases in the ethanol metabolism of *Saccharomyces cerevisiae*. *FEMS Yeast Research* 12(1): 33–47.
- Suharsono, Nurandinie D R, Tjahjoleksono A. (2020) Analysis of mutant of potato (*Solanum tuberosum* L.) cultivar Kennebec. *IOP Conference Series: Earth and Environmental Science*. <https://doi.org/10.1088/1755-1315/457/1/012074>.
- Veses V and Gow N A R. (2009). Pseudohypha budding patterns of *Candida albicans*. *Medical Mycology* 47(3): 268–275.
- Vidal A K F, Barbé T D, Daher R F, Filho J E A, Lima R S N, Freitas R S, Rossi D A *et al.* (2017). Production potential and chemical composition of elephant grass (*Pennisetum purpureum* Schum.) at different ages for energy purposes. *African Journal of Biotechnology* 16(25): 1428–1433.
- Vilela L D F, Araujo V P G D, Paredes R D S, Bon E P D S, Torres F A G, Neves B C and Eleutherio E C A. (2015). Enhanced xylose fermentation and ethanol production by engineered *Saccharomyces cerevisiae* strain. *AMB Express* 5(16): 1–7. <https://doi.org/10.1186/s13568-015-0102-y>
- Vilela L D F, Mello V M D, Reis V C B, Bon E P D S, Torres F A G, Neves B C and Eleutherio E C A. (2013). Functional expression of *Burkholderia cenocepacia* xylose isomerase in yeast increase ethanol production from a glucose-xylose blend. *Bioresource Technology*. 128(2013): 792–796.
- Walker G M and Stewart G G. (2016). *Saccharomyces cerevisiae* in the production of fermented beverages. *Beverages* 2(30): 1–12. <https://doi.org/10.3390/beverages2040030>
- Wallace-Salinas V and Gorwa-Grauslund M F. (2013). Adaptive evolution of an industrial strain of *Saccharomyces cerevisiae* for combined tolerance to inhibitors and temperature. *Biotechnology for Biofuels* 6(151): 1–9. <https://doi.org/10.1186/1754-6834-6-151>
- Young A I, Benonisdottir S, Przeworski M and Kong A. (2019). Deconstructing the sources of genotype-phenotype associations in humans. *Science* 365(6460): 1396–1400.
- Zabed H, Sahu J N, Boyce A N and Faruq G. (2016). Fuel ethanol production from lignocellulosic biomass: An overview on feedstocks and technological approaches. *Renewable and Sustainable Energy Reviews* 66: 751–774. <https://doi.org/10.1016/j.rser.2016.08.038>


Zhang W and Geng A. (2012). Improved ethanol production by a xylose-fermenting recombinant yeast strain constructed through a modified genome shuffling method. *Biotechnology for Biofuels* 5(46): 1–11. <https://doi.org/10.1186/1754-6834-5-46>

SUPPLEMENTARY MATERIAL

1. Standard Curve


- Ethanol Standard Curve

Concentration (%) [nRIU*s]16	Area
00	7182290.0
8	3624610.000
4	1835570.000
0.5	206658.000




- **Glucose Standard Curve**

Concentration (g/L) [nRIU*s]50	Area
	5755390.0
00	
40	4684250.000
30	3598550.000
20	2503300.000
10	1528590.000




- **Xylose Standard Curve**

Concentration (g/L)	Area [nRIU*s]
50	5571150.000
40	4499880.000
30	3715680.000
20	2391470.000
10	1366930.000




- **Glycerol Standard Curve**

Concentration (g/L)	Area [nRIU*s]
2	242931.000
1	119783.000
0.5	63188.300
0.25	29738.200
0.1	11820.800




- **Lactate Standard Curve**

Concentration (%)	Area [nRIU*s]
0.125	138300.000
0.25	259592.000
0.75	648493.000
1	953231.000




- **Acetate Standard Curve**

Concentration (%)	Area [nRIU*s]
0.125	112977.000
0.25	206550.000
0.5	387713.000
0.75	527406.000
1	688931.000



2. Mutant Growth on YPG and YPX media

Wild types and Fusant	Media	
	YPG*	YPX*
<i>Saccharomyces cerevisiae</i> (DBY1)	No Dilution 	No Dilution 
	<i>Candida tropicalis</i> (KBKTI 10-5.1)	No Dilution 
Mutant (Fcs)	No Dilution 	Dilution 1:10 

* Yeast cultures are spread as much as 100 L and incubated for 3 days at 30°C

3. Concentration of xylose and glucose elephant grass hydrolyzate

	Acid Hydrolyzate				Alkaline hydrolyzate			
	1%		3%		1%		3%	
	Xylose (g/L)	Glucose (g/L)	Xylose (g/L)	Glucose (g/L)	Xylose (g/L)	Glucose (g/L)	Xylose (g/L)	Glucose (g/L)
Lignocellulose of Elephant Grass (1 month old)	3.03	0.95	3.08	1.11	0.59	1.06	0.47	1.30

4.

4. Performance of ethanol production the parental and mutant yeast on YPXG media

Isolate	Ethanol (%)				Glucose (g/L)				Xylose (g/L)				Glycerol (g/L)				Lactate (g/L)				Acetate (g/L)				OD600						
	Fermentation Time (h)																														
	0	24	48	72	0	24	48	72	0	24	48	72	0	24	48	72	0	24	48	72	0	24	48	72	0	24	48	72	0	24	48
Sc	0.0	2.31	2.36	2.37	50	0.00	0.01	0.02	20	18.47	17.57	17.55	0.0	1.81	1.68	1.76	0.0	0.81	1.02	1.16	0.0	0.00	0.07	0.06	0.4	3.40	4.70	3.80			
Ct	0.0	2.24	2.24	2.29	50	0.02	0.01	0.01	20	17.00	16.29	16.23	0.0	1.47	1.48	1.38	0.0	3.18	3.58	3.38	0.0	0.06	0.06	0.01	0.4	3.90	4.70	4.37			
Fcs1	0.0	2.15	2.27	2.27	50	1.47	0.02	0.02	20	16.62	16.12	15.18	0.0	1.60	1.68	1.61	0.0	2.80	4.41	5.15	0.0	0.05	0.10	0.11	0.4	3.37	4.80	4.00			
Fcs4	0.0	2.20	2.22	2.27	50	0.00	0.02	0.02	20	16.69	16.16	15.64	0.0	1.30	1.41	1.36	0.0	2.77	3.63	3.20	0.0	0.12	0.11	0.09	0.4	3.87	5.20	4.80			

5. Performance of ethanol production the parental and mutant yeast on acid hydrolyzate

Isolate	Ethanol (%)				Glucose (g/L)				Xylose (g/L)				Glycerol (g/L)				Lactate (g/L)				Acetate (g/L)				OD600						
	Fermentation Time (h)																														
	0	4	8	24	0	4	8	24	0	4	8	24	0	4	8	24	0	4	8	24	0	4	8	24	0	4	8	24	0	4	8
Sc	0.0	0.74	0.74	0.79	1.11	0.19	0.00	0.00	3.08	2.99	2.98	2.92	0.0	0.30	0.35	0.37	0.0	0.02	0.01	0.01	0.0	0.00	0.00	0.00	0.37	0.60	1.00	1.20			
Ct	0.0	0.86	0.86	0.87	1.11	0.01	0.00	0.00	3.08	2.41	1.74	1.27	0.0	0.07	0.09	0.25	0.0	0.02	0.01	0.11	0.0	0.00	0.00	0.00	0.50	0.69	1.63	2.16			
Fcs1	0.0	0.94	0.98	1.00	1.11	0.02	0.00	0.00	3.08	2.41	2.07	1.27	0.0	0.08	0.11	0.31	0.0	0.02	0.01	0.11	0.0	0.00	0.00	0.00	0.46	0.5	1.49	2.57			
Fcs4	0.0	0.95	1.04	1.06	1.11	0.01	0.00	0.00	3.08	2.47	2.13	1.18	0.0	0.11	0.12	0.29	0.0	0.02	0.01	0.2	0.0	0.00	0.00	0.00	0.48	0.77	1.57	2.60			

6. Performance of ethanol production the parental and mutant yeast on alkaline hydrolyzate

Isolate	Ethanol (%)				Glucose (g/L)				Xylose (g/L)				Glycerol (g/L)				Lactate (g/L)				Acetate (g/L)				OD600						
	Fermentation Time (h)																														
	0	4	8	24	0	4	8	24	0	4	8	24	0	4	8	24	0	4	8	24	0	4	8	24	0	4	8	24	0	4	8
Sc	0.0	0.76	0.76	0.78	1.30	0.15	0.14	0.00	0.47	0.47	0.44	0.41	0.0	0.31	0.32	0.35	0.0	0.02	0.02	0.03	0.0	0.00	0.00	0.00	0.37	0.39	0.74	0.89			
Ct	0.0	0.69	0.74	0.74	1.30	0.18	0.14	0.00	0.47	0.33	0.18	0.00	0.0	0.24	0.26	0.04	0.0	0.01	0.02	0.02	0.0	0.00	0.00	0.00	0.50	0.74	0.82	1.51			
Fcs1	0.0	0.74	0.75	0.75	1.30	0.19	0.17	0.00	0.47	0.33	0.28	0.00	0.0	0.26	0.27	0.06	0.0	0.01	0.02	0.02	0.0	0.00	0.00	0.00	0.46	0.46	0.97	1.78			
Fcs4	0.0	0.74	0.74	0.75	1.30	0.14	0.10	0.00	0.47	0.30	0.18	0.00	0.0	0.26	0.27	0.07	0.0	0.01	0.02	0.03	0.0	0.00	0.00	0.00	0.48	0.54	0.92	1.73			