



Histopathological Alterations in Gills, Liver and Kidney of African Catfish (*Clarias gariepinus*, Burchell 1822) Exposed to *Melaleuca cajuputi* Extract

Authors:

Marina Hassan*, Anuar Abdalah Nagi Melad, Mohd Ihwan Zakariah and Nor Asma Husna Yusoff

***Correspondence:** marina@umt.edu.my

Submitted: 23 December 2021; **Accepted:** 3 October 2022; **Early view:** 11 November 2022

To cite this article: Marina Hassan, Anuar Abdalah Nagi Melad, Mohd Ihwan Zakariah and Nor Asma Husna Yusoff (in press). Histopathological alterations in gills, liver and kidney of African catfish (*Clarias gariepinus*, Burchell 1822) exposed to *Melaleuca cajuputi* extract. *Tropical Life Sciences Research*.

Highlights

- The severity of gills, liver and kidney of African catfish were proportional to the increase of MCLE concentrations.
- Degeneration alterations were notable in gills, liver and kidney of African catfish when exposed to MCLE concentration over than the LC₅₀ value.
- It was recommended that MCLE exposure be less than the LC₅₀ value to lessen the histopathological alterations in the gills, liver, and kidneys of African catfish.

Accepted author version posted online: 11 November 2022

Histopathological Alterations in Gills, Liver and Kidney of African Catfish (*Clarias gariepinus*, Burchell 1822) Exposed to *Melaleuca cajuputi* Extract

¹Marina Hassan*, ^{1,2}Anuar Abdalah Nagi Melad, ¹Mohd Ihwan Zakariah and ¹Nor Asma Husna Yusoff

¹Higher Institution Centre of Excellence (HICoE), Institute of Tropical Aquaculture and Fisheries, Universiti Malaysia Terengganu, 21300 Kuala Nerus, Terengganu, Malaysia

²Department of Biological Science, Faculty of Science, Azzaytuna University, Tarhunah, Libya

*Corresponding author: marina@gmail.com; marina@umt.edu.my

Running head: Histopathological effect on African catfish organs exposed to *M. cajuputi*

Abstract: This study evaluated the histopathological changes in the gill, liver and kidney of African catfish (*Clarias gariepinus*) intoxicated with a sub-lethal dose of *Melaleuca cajuputi* leaves extract (MCLE) for 96 hours. The acute toxicity test has been determined previously with a value of 96-h LC₅₀ = 127 mg/L, hence the selection of sub-lethal ranges from 60 to 160 mg/L of MCLE. Degenerative alterations were prominent in all tested organs, particularly after exposure to a high concentration of MCLE. Gill exhibited haemorrhage, epithelial lifting, lamellar disorganization, and necrosis after exposure to a high MCLE concentration. Alterations in the liver include congestion, hydropic degeneration, and vacuolation, whereas lesions in the kidney were pyknosis, vacuolation, hydropic degeneration, and tubular necrosis. The obtained data showed that the organs experienced severe changes proportional to the increase in MCLE concentration. In addition, fish exposed to higher concentrations than the LC₅₀ value experienced irreversible lesions. The present study suggests that the use of MCLE below the LC₅₀ is recommended to avoid severe alterations to organs, particularly in African catfish. This study demonstrated that the use of MCLE above the LC₅₀ promotes severe damage to the gills, liver, and kidney of African catfish. However, further investigations are needed to define the causing-mechanisms underlying these effects.

Keywords: Gelam, Catfish, Histopathology Alterations, Sub-Lethal

Abstrak: Kajian ini menilai perubahan histopatologi dalam insang, hati, dan buah pinggang ikan keli Afrika (*Clarias gariepinus*) yang ditoksikkan dengan dos sub-lethal ekstrak daun *Melaleuca cajuputi* (MCLE) selama 96 jam. Ujian ketoksikan akut telah ditentukan sebelum ini dengan nilai 96-jam LC₅₀ = 127 mg/L, oleh itu pemilihan julat sub-lethal dari 60 hingga 160 mg/L MCLE. Perubahan degeneratif adalah ketara dalam semua organ yang diuji, terutamanya selepas terdedah kepada kepekatan MCLE yang tinggi. Insang menunjukkan pendarahan, pengangkatan epitelium, kekacauan lamellar, dan nekrosis selepas terdedah kepada kepekatan MCLE yang tinggi. Perubahan dalam hati termasuk kesesakan, degenerasi hidropik, dan vakuolasi, manakala luka pada buah pinggang adalah piknosis, vakuolasi, degenerasi hidropik, dan nekrosis tiub. Data yang diperolehi menunjukkan bahawa organ-organ mengalami perubahan teruk yang berkadar dengan peningkatan kepekatan MCLE. Selain itu, ikan yang terdedah kepada kepekatan yang lebih tinggi daripada nilai LC₅₀ mengalami luka yang tidak dapat dipulihkan. Kajian ini mencadangkan bahawa penggunaan MCLE di bawah nilai LC₅₀ disyorkan untuk mengelakkan perubahan teruk pada organ, terutamanya pada ikan keli Afrika. Kajian ini menunjukkan bahawa penggunaan MCLE di atas nilai LC₅₀ menggalakkan kerosakan teruk pada insang, hati, dan buah pinggang ikan keli Afrika. Walau bagaimanapun, penyiasatan lanjut diperlukan untuk menentukan mekanisme punca yang menjadi penyebab kepada kesan ini.

Kata kunci: Gelam, Ikan Keli, Perubahan Histopathologi, Sub-Letal

INTRODUCTION

Aquaculture is one of the fastest growing sectors in the food industry, accounting for more than 20 % of all animal protein exported to low-income countries worldwide (Tacon 2020). Among them, fish farming accounts for nearly 40 % (54.3 million tons), followed by mollusks and crustaceans at 17.7 million tonnes and 9.4 million tons, respectively (FAO 2020). However, intensive aquaculture with high stocking capacity promotes the spread of pathogenic microorganisms, resulting in massive economic losses (Valenzuela-Gutiérrez *et al.* 2021). Previously, several chemical antibiotics such as formalin, trichlorfon, and mebendazole were used to control this situation (Liu *et al.* 2021), but their overuse resulted in antibiotic accumulation, which might pose health risks to the consumer (Aisiah *et al.* 2020). Furthermore, long-term use of these antibiotics may result in bacterial disease resistance and a lack of efficacy in some control

agents (Zhang *et al.* 2014). To avoid the use of chemotherapeutical treatments, fish need to improve their immunity to combat the dispersion of diseases and pathogens. The use of plant extracts as immunostimulants in aquatic organisms has been reported worldwide through improving growth and body defence mechanisms (Sahimi *et al.* 2022; Doan *et al.* 2022; Rashidian *et al.* 2021; Nafiqoh *et al.* 2020; Mohammadi *et al.* 2020). Therefore, the use of plant extracts as an alternative is promising as a potential strategy to improve fish immunity to combat diseases and parasites due to their advantages of being environmentally friendly and biodegradable (Liu *et al.* 2021).

Melaleuca cajuputi is one of the native plants that are commonly found throughout Southeast Asia and Australasia (Nagi *et al.* 2016). Major phytochemical compounds such as terpenes, alkaloids, phenolics, and flavonoids were previously reported in *M. cajuputi* extract which contribute to a broad-spectrum of biological activities (Chaudhari *et al.* 2022; Sahimi *et al.* 2022; Septiana *et al.* 2020; Bua *et al.* 2018; Chan and Haron 2016; Al-Abd *et al.* 2015). Some of them demonstrated their toxicity through acute toxicity tests (Jayakuamar *et al.* 2020; Noor *et al.* 2020; Bakar 2020; Roszaini *et al.* 2013), but not all of the activity has been effectively characterised *in vivo* to determine their safety effect at a sub-lethal dose. There are few reports of histopathological changes following the use of plant extracts on aquatic organisms (Meneses *et al.* 2020; Karataş *et al.* 2020; Meneses *et al.* 2018; Shivappa *et al.* 2015), but there is a paucity in the *M. cajuputi* plant extract. Therefore, this study was performed to determine the histopathological changes occur in the gills, liver, and kidney of African catfish at sub-lethal doses. Earlier, an acute toxicity study of MCLE had been done on the catfish, *Clarias gariepinus*, which gave a 96-hour LC₅₀ of 127 mg/L (Nagi *et al.* 2016). The findings may be useful in determining the safety of using MCLE as a baseline for fish, particularly African catfish, as therapeutics.

MATERIALS AND METHODS

Collection and Identification of *Melaleuca cajuputi* Powell

Fresh *Melaleuca cajuputi* leaves were collected in Merang, Terengganu (5.4932°N, 102.9320°E) and sent to botanist for taxonomy identification under the voucher number of MFI 0244/22.

Preparation of *M. cajuputi* leaves extract (MCLE) and treatment solution

Leaves were washed under running tap water to remove any soil debris and dried at room temperature (27 ± 2 °C) for 14 days. Dried leaves were ground and weighed for 10 g before soaking in 80 % methanol for 24 hours with shaking (27 °C, 200 rpm). The mixture was then filtered using filter paper (Whatman, UK) and the filtrate was further concentrated using a rotary evaporator (Buchi Rotavapor-R, Fisherscience, UK) at 38 °C until the gummy-like crude extract was formed. The crude extract was freeze-dried and the powder form was stored at -20 °C until further use.

Phytochemical Compound Identification using Gas Chromatography-Mass Spectra (GC-MS) Analysis

GC-MS analysis was performed according to Al-Abd *et al.* (2015), with slight modifications. Briefly, 10 mg of MCLE sample was sonicated for 15 min in 2.5 mL of dichloromethane at 40 °C in a sealed vial. Then, 1 mL of the mixture was filtered through a 0.20 µM nylon filter, and 1 µl of the filtrate was injected splitless into the GC-MS system (Hawlett-Packard autosampler 6890 GC, Agilent, USA), equipped with a capillary column (diameter: 30 m × 0.25 mm; thickness: 0.24 µM). The column initial temperature was initially set up at 40 °C for 3 min, and the final temperature was at 280 °C for 3 min. Helium was used as a carrier gas at 1 ml/min flow rate. The ion source temperature and injector were both maintained at 250 °C. The mass spectrometer was scanned over the range of 28–400 m/z amu at 1 scan⁻¹, with an ionising voltage of 70 eV. The spectra that came up were then compared with the known compounds from the NIST database library based on their retention time (RT) index of primary and secondary metabolites.

Fish Acclimatization

Clarias gariepinus with a mean body weight and total length of 20.0 ± 3.0 g and 15.0 ± 2.0 cm, respectively, were obtained from the hatchery at the Institute of Tropical Aquaculture and Fisheries, Universiti Malaysia Terengganu (UMT). Fish were placed in a tank containing 500 L of non-chlorinated water and acclimatized for two weeks under laboratory conditions. The fish were fed daily with commercial fish pellets at 3 % of their body weight. The water quality parameters, including temperature, pH, and dissolved oxygen, were measured using multiparameter

equipment (YSI 556 MPS, USA) and maintained at 26–28 °C, 5.50–6.25, and >5 mg/L, respectively. Approximately 80–90 % of the water was changed daily to keep the quality.

Experimental Procedure

The acute toxicity and behavioral studies have been performed previously by Nagi *et al.* (2016), which indicated the 96-hrs LC₅₀ value of MCLE to be at 127 mg/L. In this present study, a sub-lethal toxicity test was performed to determine the histological changes on gills, liver, and kidney of African catfish after exposure to MCLE for 96 hrs, as according to Correia *et al.* (2020), with slight modifications. Briefly, six MCLE concentration were used in this study, (i.e., 0.0, 60, 80, 100, 120, 140, and 160 mg/L), where 0.0 mg/L was served as control. Healthy acclimatized fish were randomly distributed in 18 aquaria (50 L) containing MCLE solutions, with 5 fish per treatment, and in triplicates (5 fish × 6 MCLE concentrations × 3 replicates). Water parameters including temperature, pH and dissolved oxygen were controlled at 26.28C, 5.5-6.25, and >5 mg/L, respectively using YSI multi-parameter probe (YSI Inc., USA). No feeding was performed during the experiment.

Histological Assessment of Gills, Liver and Kidney of African Catfish

At the end of experiment, fish were euthanized by rapid cooling (Rodrigues *et al.* 2019), decapitated, and the gills, liver, and kidney were carefully removed. Thereafter, the organ tissues were fixed in 10% formalin solution, followed by dehydration through six grades of alcohol separately, sectioned using a microtome (Leica, Germany) sectioning, and stained with H&E. Finally, the mounted slides were examined under a light microscope (Leica, Germany) to observe any histological changes in the tissues. Histopathological changes observed in tissue were captured using a digital camera attached with software (Leica Application Suitr V4.0, Leica, Germany) at various magnifications. The histopathological alterations were evaluated using quantitative histological assessment as adopted by Bernet *et al.* (1999), and Nero *et al.* (2006), with some modifications. Each organs were classified into four reaction patterns (rp): circulatory, degeneration, inflammatory, and structural. The score values from 1 to 6 (a) were assigned based on the percentage of each alteration, whereas the importance factors (w) ranging from 1 to 3 indicate how the alterations might affect the fish health The calculation for each category of alterations and the total pathological index were defined as:

$$\text{Org}_{\text{cat}} = \sum_{\text{alt}} (a\text{Org}_{\text{cat_alt}} \times w\text{Org}_{\text{cat_alt}})$$

$$T_{\text{org}} = \sum_{\text{cat}} \sum_{\text{alt}} (a\text{Org}_{\text{cat_alt}} \times w\text{Org}_{\text{cat_alt}}),$$

where: org = organ; cat = category; alt = alteration; a = score; w = importance factor.

Statistical Analysis

Data normality were checked prior to statistical analysis using normal probability plots. The statistical differences between the control and MCLE-exposed fish were determined using one-way analysis of variance (ANOVA), and Tukey pairwise test at $p < 0.05$ to indicate the significant difference relative to the control group (Genstat Software ver. 12.1).

RESULTS

Determination of Phytochemical Compounds in MCLE using GC-MS Analysis

A total of thirty-seven compounds were identified in MCLE as shown in Table 1. There are four major compounds were detected including 1H-Imidazole, 4,5-diphenyl- (18.60%), followed by 7-Methyl-6,8-bis(methylthio)pyrrolo[1,2]pyrazine (15.43%), 2-Isopropyl-10-methylphenanthrene (13.50%), and 9-carboxaldehyde-10-Methylantracene (11.13%) within 30 minutes of retention time (Fig. 1).

Table 1: Phytochemical compounds in MCLE identified by GC-MS analysis.

No.	RT (min)	Name of the compound	Molecular Formula	Molecular Weight	Peak Area (%)
1	5.598	1-(1-methylethyl)- 4-methyl-3-Cyclohexen-1-ol,	C ₁₀ H ₁₈ O	154.2493	0.25
2	5.655	Benzene, 1-methyl-4-(1-methylethenyl)-	C ₁₀ H ₁₂	132.2023	0.38
3	5.724	3-Cyclohexene-1-methanol, .alpha.,.alpha.,4-trimethyl-, (S)-	C ₁₀ H ₁₈ O	154.25	0.35
4	8.957	Naphthalene, 1,2,3,5,6,7,8,8a-octahydro-1,8a-dimethyl-7-(1-methylethenyl)	C ₁₅ H ₂₄	204.3511	0.19
5	9.048	Naphthalene, 1,2,3,4,4a,5,6,8a-octahydro-4a,8-dimethyl-2-(1-methylethenyl)	C ₁₅ H ₂₄	204.3511	0.26

6	9.2941	Naphthalene, 1,2,3,5,6,8a-hexahydro-4,7-dimethyl-1-(1-methylethyl)	C ₁₅ H ₂₄	204.3511	0.22
7	9.569	Cyclohexanemethanol, 4-ethenyl-.alpha.,.alpha.,4-trimethyl-3-(1-methylethenyl)	C ₁₅ H ₂₆ O	222.3663	0.52
8	9.912	1H-Cycloprop[e]azulen-7-ol, decahydro-1,1,7-trimethyl-4-methylene-, [1ar-(1a.alpha.,4a.alpha.,7.beta.,7a.beta.,7b.alpha.)]-	C ₁₅ H ₂₄ O	220.3505	2.26
9	9.981	Caryophyllene oxide	C ₁₅ H ₂₄ O	220.35	2.07
10	10.084	Guaiol	C ₁₅ H ₂₆ O	222.37	3.32
11	10.227	7-Methyl-6,8-bis(methylthio)pyrrolo[1,2]pyrazine	C ₁₀ H ₁₂ N ₂ S	224.34	15.43
12	10.398	Cycloisolongifolene, 8,9-dehydro	C ₁₅ H ₂₂	202.3352	1.66
13	10.438	2-Naphthalenemethanol, 1,2,3,4,4a,5,6,7-octahydro-.alpha.,.alpha.,4a,8-tetramethyl-, (2R-cis)	C ₁₅ H ₂₆ O	222.3663	1.96
14	10.496	6-Isopropenyl-4,8a-dimethyl-1,2,3,5,6,7,8,8a-octahydro-naphthalen-2-ol	C ₁₅ H ₂₄ O	220.3505	1.14
15	10.644	2-Naphthalenemethanol, decahydro-.alpha.,.alpha.,4a-trimethyl-8-methylene-, [2R-(2.alpha.,4a.alpha.,8a.beta.)]-	C ₁₅ H ₂₆ O	222.3663	7.25
16	10.776	1H-Cycloprop[e]azulen-4-ol, decahydro-1,1,4,7-tetramethyl-, [1ar-(1a.alpha.,4.beta.,4a.beta.,7.alpha.,7a.beta.,7b.alpha.)]-	C ₁₅ H ₂₆ O	222.3663	1.42
17	10.810	Humulen-(v1)	C ₁₅ H ₂₄	204.3510	0.97
18	10.956	7R,8R-8-Hydroxy-4-isopropylidene-7-methylbicyclo[5.3.1]undec-1-ene	C ₁₅ H ₂₄ O		0.70
19	11.148	2-Acetyl-5-chloro-3-methylbenzo(b)thiophene	C ₁₁ H ₉ ClO S	224.7066	5.45
20	11.280	Bicyclo[4.3.0]nonane, 7-methylene-2,4,4-trimethyl-2-vinyl-	C ₁₅ H ₂₄	204.3510	0.81
21	11.308	6-Isopropenyl-4,8a-dimethyl-1,2,3,5,6,7,8,8a-octahydro-naphthalen-2-ol	C ₁₅ H ₂₄	204.3510	0.77
22	11.743	1-(3-Chloromethyl-2,4,6-trimethylphenyl)ethanone	C ₁₂ H ₁₅ ClO	210.6999	1.43
23	11.783	p-Isopropylphenetole	C ₁₁ H ₁₆ O	164.2441	0.21
24	12.064	2',3',4',5',6'-Pentafluoroacetophenone	C ₈ H ₃ F ₅ O	210.1	0.43
25	12.149	2-Naphthalenemethanol, 1,2,3,4,4a,5,6,8a-octahydro-.alpha.,.alpha.,4a,8-tetramethyl-, [2R-(2.alpha.,4a.alpha.,8a.beta.)]-	C ₁₅ H ₂₆ O	222.3663	0.57
26	12.727	4H-1-Benzopyran-4-one, 5-hydroxy-7-methoxy-2-methyl-	C ₁₁ H ₁₀ O ₄	206.19	0.59
27	12.956	Hexadecanoic acid, methyl ester	C ₁₇ H ₃₄ O ₂	270.4507	0.29
28	13.299	Hexadecanoic acid	C ₁₆ H ₃₂ O ₂	256.4241	0.42
29	13.591	9-carboxaldehyde-10-Methylanthracene	C ₁₆ H ₁₂ O	220.27	11.13
30	13.643	9H-Fluoren-9-ol, 9-butyl-	C ₁₇ H ₁₈ O	238.3242	3.04
31	13.780	1H-Imidazole, 4,5-diphenyl-	C ₁₅ H ₁₂ N ₂	220.2692	18.60

32	13.940	2-Isopropyl-10-methylphenanthrene	C ₁₈ H ₁₈	234.34	13.50
33	14.495	3,7,11,15-Tetramethyl-2-hexadecen-1-ol			0.50
34	14.695	9,12,15-Octadecatrien-1-ol	C ₁₈ H ₃₂ O	264.4461	0.42
35	17.677	1H-Indole, 5-methyl-2-phenyl-	C ₁₅ H ₁₃ N	207.2704	0.22
36	17.763	5-Methyl-2-phenylindolizine			0.27
37	26.088	Stigmasterol, 22,23-dihydro-	C ₂₉ H ₅₀ O	414.7067	1.02




Figure 1: Total ion chromatogram of MCLE by GC-MS analysis.

Histopathological assessment

Overall, the gills, liver, and kidney of African catfish showed significant pathological alterations when exposed to a higher concentration of MCLE ($p < 0.05$) than the control. All organs showed pathological indices associated with the elevation of MCLE concentrations, particularly starting at 120 to 160 mg/L. Degenerative alterations predominated ($p < 0.05$) in all the tested organs of MCLE-treated fish compared to control (Figs. 2A, 2B, 2C). Concerning to gills and liver, necrosis was recorded at higher MCLE concentrations (120 to 160 mg/L), whereas started later at 140 to 160 mg/L in the kidney. However, nuclear alterations (e.g. pyknosis) and hydropic degeneration were already observed in the kidney of fish exposed to 120 mg/L. In terms of circulatory changes, sinusoids congested with red blood cells and cytoplasmic vacuolation were observed in the liver at lower exposures of MCLE. Among the reported circulatory changes, haemorrhage was the most detected in fish kidney, even increasing the lowest exposure of MCLE concentration.




Figure 2: A Pathological indices of *C. gariepinus* gills at different concentration of MCLE. (C) Control; (T1) 60 mg/L; (T2) 80 mg/L; (T3) 100 mg/L; (T4) 120 mg/L; (T5) 140 mg/L; (T6) 160 mg/L. (*): have significant different at $p < 0.05$




Figure 2: B Pathological indices of *C. gariepinus* liver at different concentration of MCLE. (C) Control; (T1) 60 mg/L; (T2) 80 mg/L; (T3) 100 mg/L; (T4) 120 mg/L; (T5) 140 mg/L; (T6) 160 mg/L. (*): have significant different at $p < 0.05$









Figure 2: C Pathological indices of *C. gariepinus* kidney at different concentration of MCLE. (C) Control; (T1) 60 mg/L; (T2) 80 mg/L; (T3) 100 mg/L; (T4) 120 mg/L; (T5) 140 mg/L; (T6) 160 mg/L. (*): have significant different at $p < 0.05$

No histological changes were observed in the gills, liver, and kidney of non-treated fish (Fig 3A). However, several changes were observed in MCLE-treated fish with the severity increasing as the plant extract concentration increased. In the 60 mg/L MCLE (Fig 3B), a minor degree of oedema, hydropic degeneration of the epithelial cells, and lamella fusion were observed. High tissue alteration was observed in 80 mg/L MCLE (Fig 3C), whereas lesion was observed in 100 mg/L MCLE, including epithelial lifting, congestion, lamellar epithelial cell degeneration, and oedema (Fig 3D). Severe lesions including haemorrhage, lamella fusion, oedema, and necrosis were observed in 120 mg/L MCLE (Fig 3E). Severe degenerative and necrotic changes in gill epithelia were detected in gills exposed to 140 and 160 mg/L MCLE (Fig 3F, 3G), with huge lamellar disorganisation and aneurisms found only in 160 mg/L MCLE (Fig 3G).












Figure 3: **A** Normal gills of *C. gariepinus* (0 mg/L MCLE). **B** 60 mg/L of MCLE with 1. Oedema, 2. Hydropic degeneration of the epithelial cells. **C** 80 mg/L MCLE with 1. Epithelial lifting, 2. Vines congestion, 3. Oedema, 4. Hydropic degeneration of the epithelial cells. **D** 100 mg/L MCLE with 1. Epithelial lifting, 2. Vines congestion, 3. Epithelial cells degeneration, 4. Oedema. **E** 120 mg/L MCLE with 1. Haemorrhage, 2. Lamellar epithelial cells degeneration, 3. Oedema, 4. Necrosis. **F** 140 mg/L MCLE with 1. Epithelial lifting of secondary lamellae, 2. Necrosis, 3. Hydropic degeneration of the epithelial cells. **G** 160 mg/L MCLE with 1. Lamellar disorganization, 2. Haemorrhage, 3. Necrosis, 4. Oedema. H&E stain, 400x total magnifications.

No lesions was observed in non-treated fish liver (Fig 4A). Meanwhile, alterations were observed in liver tissues at 60 mg/L MCLE, including congestion of sinusoids and vacuolation (Fig 4B). A slight changes like plasma alteration, degeneration, and leukocyte infiltration in fish exposed to 80 mg/L MCLE (Fig 4C). More alteration was found in higher MCLE concentration such as congested of central vein with poor hepatic cord structure (Fig 4D) and hepatocytes focal necrosis (Fig 4E), and even more severe in the higher MCLE concentration (Fig 4F). For the kidney, a normal appearance was observed in the untreated kidney tissues (Fig 5A). However, a slight change was observed with the increased plant concentration (Fig 5B), including tubular congestion with red blood cells and hydropic degeneration (Fig 5C), vacuolation, hydropic degeneration of tubular cells (Fig 5D), and severe hydropic degeneration of tubular cells, haemorrhage, and tubular necrosis (Fig 5E, 5F).







Figure 4: **A** Normal liver of *C. gariepinus* (0 mg/L MCLE) with 1. Hepatocytes, 2. Hepatic portal vein, 3. Hepatic artery, 4. Bile duct. **B** 60mg/L MCLE with 1. Sinusoid congested with red blood cells, 2. Cytoplasmatic vacuolation. **C** 80 mg/L MCLE with 1. Central vein congested with red blood cells, 2. Infiltration of leukocytes, 3. Vacuolation, 4. Cloudy degeneration, 5. Aggregation of inflammatory cell between hepatocytes. **D** 100 mg/L MCLE with 1. Structure alteration. **E** 120 mg/L MCLE with 1. Focal necrosis of hepatocytes, 2. Hydropic degeneration. **F** 140 mg/L MCLE with 1. Vacuolation, 2. Degeneration, 3. Necrosis, 4. Increase in sinusoidal spaces and congested with red blood cells. **G** 160 mg/L MCLE with severely necrotic liver tissues, vacuolation of hepatocytes, hydropic degeneration of hepatocytes and poor hepatic cord structural. H&E stain, 400x total magnifications.






Figure 5: **A** Normal kidney of *C. gariepinus* (0 mg/L MCLE) with 1. Glomerulus, 2. Bowman's space well defined, 3. Proximal tubule, 4. Distal tubule, 5 Bowman's capsule, 6. Hematopoietic tissue. **B** 60 mg/L MCLE with 1. Haemorrhage. **C** 80 mg/L MCLE with 1. Haemorrhage, 2. Tubular degeneration. **D** 100 mg/L MCLE with 1. Shrinkage of glomerulus, 2. Vacuolation and hydropic degeneration. **E** 120 mg/L MCLE with 1. Nuclear alterations, 2. Vacuolation and hydropic degeneration. **F** 140 mg/L MCLE with 1. Haemorrhage, 2. Tubular necrosis. **G** 160 mg/L MCLE with 1. Severely necrotic kidney glomerulus, 2. Vacuolation, 3. Tubular degeneration, 4. Dilatation

of glomerulus capillaries, 5. Hematopoietic tissue with red blood cells and poor tubular structural. H&E stain, 400x total magnifications.

DISCUSSION

The histopathological evaluation of tissue and cells caused by toxicants is crucial to determining the health condition of organisms. In this present study, *M. cajuputi* leaves extract (MCLE) was exposed for 96 hrs to African catfish at sub-lethal doses, and the histological alterations in the gills, liver, and kidney were observed. In the GC-MS analysis, 37 compounds were detected in MCLE, with terpenes and flavonoids being the major compounds. Terpenes such as caryophyllene oxide (2.07 %), 1H-Imidazole, 4,5-diphenyl-18.60%, 2-isopropyl-10-methylphenanthrene (13.50 %), and stigmasterol (1.02 %) are among the higher peaks found in MCLE, which have also been reported previously (Noor *et al.* 2021; Białoń *et al.* 2019; Ukit *et al.* 2019), however, with a slight difference in quantity, which might be influenced by several geographical factors including method of extraction, harvesting period, part of the plant used and more (Kumari *et al.* 2014). These compounds may act synergistically or in a single compound to serve a variety of biological properties such as antibacterial (Gonzalez *et al.* 2021; Moghrovyan *et al.* 2019), anti-inflammatory (Al-Khayri *et al.* 2022; Prado-Audelo *et al.* 2021), antioxidant (Ivanović *et al.* 2020; Amri and Hossain 2018), antifungal (Salomé-Abarca *et al.* 2020; de Lima *et al.* 2022), and many more.

An acute toxicity test of MCLE on African catfish has been conducted previously by Nagi *et al.* (2016), which resulted in a 96-hour LC₅₀ of 127 mg/L. Following that, a sub-lethal toxicity test was performed in this current study to determine the histopathological effects on the gills, liver, and kidney of African catfish during the exposed time. Gills are a good indicator of water quality because of their large surface area and contacts with the external environment (Strzyzewska *et al.* 2016). The harmful chemical and biological substances in the water surroundings may influence the gills' morphological changes and impairment of their functions. The results of this study revealed that histopathological changes such as epithelial lifting, oedema, and hydropic degeneration were observed in fish gills exposed to lower concentrations of MCLE, but the effects increased proportionally with MCLE concentrations, such as lamellar disorganisation and necrosis. It can be speculated that these significant alterations may be attributed to the high content of phytochemical compounds in MCLE. Although mucous secretion aids in the prevention of toxicants passing to the gill epithelium, these consequences may disturb

the gas exchange process and reduce respiration (Nero *et al.* 2006). Similar findings were reported by Pal *et al.* (2012) after exposure of chlorpyrifos to common carp, *Cyprinus carpio*. The common alterations observed in all concentrations were epithelial lifting and hydropic degeneration, which increased proportionally with toxicant concentrations. Similar alterations were found in *Sparus aurata* following acute and chronic exposure to erythromycin and oxytetracycline (Rodrigues *et al.* 2019). These alterations may be induced by severe oedema, leading to a partial or complete fusion of the gill filament (Yancheva *et al.* 2016).

The liver is a primary organ for detoxification of toxicants, storage of nutrients, metabolism, and synthesis of enzymes; it therefore becomes highly vulnerable to toxicants (Rodrigues *et al.* 2019). High exposure to MCLE resulted in focal necrosis, vacuolation, increase in sinusoidal spaces, and poor hepatic cord structure. Vacuolation can happen due to biochemical disturbances leading to slightly glycogen flocculent and single/multiple lipid accumulation in hepatocyte cytoplasm (Yancheva *et al.* 2016). Similar observations were found in fish exposed to toxicants (Nunes *et al.* 2015; Rodrigues *et al.* 2017). Liver necrosis is primarily induced by oxidative stress generated by toxicants (Verma *et al.* 2007) and was only found in higher exposures of MCLE. It may lead to irreversible liver damage like pyknotic nucleus with a long period of exposure due to nucleus hyperactivity (Pal *et al.* 2012).

A significant pathological index in the kidney was mainly attributed to degenerative alterations such as tubular degeneration coupled with nuclear alterations, which indicates that the exposure of MCLE at 100 to 160 mg/L was toxic to fish even at a shorter time. Tubule degeneration and necrosis were also reported by Correia *et al.* (2020) after acute effects of cerium dioxide nanoparticles on *Oncorhynchus mykiss*. In addition, a significant circulatory disturbance was observed in the kidney, including haemorrhage and shrinkage of the glomerulus, suggesting that MCLE affects cell permeability and causes disturbance to fluid transport in and out of cells (Hadi and Alwan 2012). Finding was contradict with Shivappa *et al.* (2015) where they discovered no severe changes was observed in liver and heart tissues of goldfish and clownfish after exposure to *M. cajuputi* extract product, *Melafix*.

CONCLUSION

During short-term application, the exposure of MCLE at acute levels showed several mild and moderate histological alterations in gills, liver, and kidney of African catfish, possibly due to effect of phytochemical compounds contain in extract. Additionally, the effects were dose-dependent,

and using high MCLE concentrations, especially above the LC₅₀ value (i.e.120 to 160 mg/L), severely damages the organs' tissues and structural integrity, potentially leading to organ malfunction. To ascertain their effectiveness, more research in other aquatic animal species may prove useful, as different animals may behave differently when exposed to plant extracts.

ACKNOWLEDGEMENTS

The research was supported by the Ministry of Higher Education, Malaysia, under the Exploratory Research Grant Scheme (ERGS) (Vot. No. 55053), and the Ministry of Higher Education, Malaysia, under the Higher Institute Centre of Excellence (HICoE) program (Vot. No. 63933 & 56051).

AUTHORS CONTRIBUTIONS

Marina Hassan is the leader who initiated the project, monitored the authors during the experiment and data analysis, and was greatly contributed in writing the manuscript. Anur Abdalah Nagi Melad conducted all the experiments, data analysis and writing the manuscript. Meanwhile, Mohd Ihwan Zakariah and Nor Asma Husna Yusoff involved in data analysis and writing the manuscript.

REFERENCES

- Aisiah S, Prajitno A, Maftuch M and Yuniarti A. (2020). Effect of *Nauclea subdita* (Korth.) Steud. leaf extract on haematological and histopathological changes in liver and kidney of striped catfish infected by *Aeromonas hydrophila*. *Veterinary World* 13: 2231-0916. www.veterinaryworld.org/Vol.13/January-2020/7.pdf
- Al-Abd N, Nor Z M, Mansor M, Azhar F, Hasan M S and Kassim M. (2015). Antioxidant, antibacterial activity, and phytochemical characterization of *Melaleuca cajuputi* extract. *BMC Complementary and Alternative Medicine* 15: 385. <http://doi.org/10.1186/s12906-015-0914-y>
- Al-Khayri J M, Sahana G R, Nagella P, Joseph B V, Alessa F M and Al-Mssallem M Q. (2022). Flavonoids as potential anti-inflammatoryMolecules: a review. *Molecules* 27: 2901. <https://doi.org/10.3390/molecules27092901>

- Amri F S and Hossain M A. (2018). Comparison of total phenols, flavonoids and antioxidant potential of local and imported ripe bananas. *Egyptian Journal of Basic and Applied Sciences* 5(4): 245–251. <http://doi.org/10.1016/j.ejbas.2018.09.002>
- Bakar A A. (2020). Bioactivity evaluation of *Melaleuca cajuputi* (Myrtales: Myrtaceae) crude extracts against *Aedes* mosquito. *Tropical Agricultural Science* 43(3): 303–313.
- Bernet D, Schmidt H, Meier W, Burkhardt-Holm P and Wahli T. (1999). Histopathology in fish: proposal for a protocol to assess aquatic pollution. *Journal of Fish Diseases* 22(1): 25–34.
- Białoń M, Krzyśko-Łupicka T, Nowakowska-Bogdan E and Wieczorek P P. (2019). Chemical composition of two different lavender essential oils and their effect on facial skin microbiota. *Molecules* 24(18): 3270. <http://doi.org/10.3390/molecules24183270>
- Bua A, Malicotti P, Donadu M G, Usai D, Le L S, Tran T T T, Ngo V Q T, Marchetti M, Usai M, Cappuccinelli P and Zanetti S. (2018). ‘*In vitro*’ activity of *Melaleuca cajuputi* against mycobacterial species. *Natural Product Research*. <https://doi.org/10.1080/14786419.2018.1509335>
- Chan B K and Haron H. (2016). Insights into Putative Health Implications of Gelam (*Melaleuca cajuputi*) Honey: Evidence from In-Vivo and In-Vitro Studies. *Medical Sciences* 4(3). <http://doi.org/10.3390/medsci4010003>
- Chaudhari A K, Singh V K, Das S, Kujur A, Deepika and Dubey N K. (2022). Unveiling the cellular and molecular mode of action of *Melaleuca cajuputi* Powell. essential oil against aflatoxigenic strains of *Aspergillus flavu* isolated from stored maize samples. *Food Control*, 138: 109000. <https://doi.org/10.1016/j.foodcont.2022.109000>
- Correia A T, Rodrigues S, Ferreira-Martins D, Nunes A C, Ribeiro M I and Antunes S C. (2020). Multi-biomarker approach to assess the acute effects of cerium dioxide nanoparticles in gills, liver and kidney of *Oncorhynchus mykiss*. *Comparative Biochemistry and Physiology, Part C* 238: 108842. <https://doi.org/10.1016/j.cbpc.2020.108842>
- de Lima L F *et al.* (2022). Anti-candida properties of *Gossypium hirsutum* L.: enhancement of fungal growth, biofilm production and antifungal resistance. *Pharmaceutics* 14: 698. <https://doi.org/10.3390/pharmaceutics14040698>
- Doan H V, Lumsangkul C, Sringarm K, Hoseinifar S H, Dawood M A O, Haron E E, Harikrishnan R, Jaturasitha S and Paolucci M. (2022). Impacts of Amla (*Phyllanthus emblica*) fruit extract on growth, skin mucosal and serum immunities, and disease resistance of Nile tilapia (*Oreochromis niloticus*) raised under biofloc system. *Aquaculture Reports* 22: 100953. <https://doi.org/10.1016/j.aqrep.2021.100953>

- FAO. (2020). *The state of the world fisheries and aquaculture 2020. Sustainability in action*. Food and Agriculture Organization of the United Nations.
- González A, Casado J and Lanas A. (2021). Fighting the antibiotic crisis: flavonoids as promising antibacterial drugs against *Helicobacter pylori* Infection. *Frontiers in Cellular and Infection Microbiology* 11: 709749. <http://doi.org/10.3389/fcimb.2021.709749>
- Hadi A A and Alwan S F. (2012). Histological changes in gills, liver and kidney of freshwater fish, *Tilapia zillii*, exposed to aluminium. *International Journal of Pharmaceutical Life Sciences* 3(11): 2071–2081.
- Ivanović S, Avramović N, Dojčinović B, Trifunović S, Novaković M, Tešević V and Mandić B. (2020). Chemical composition, total phenols and flavonoids contents and antioxidant activity as nutritive potential of roasted hazelnut skins (*Corylus avellana* L.). *Foods* 9: 430. <http://doi.org/10.3390/foods9040430>
- Jayakumar M, Ramachandran M, Krisnaveni T and Nattudurai G. (2020). Toxicity and biochemical effects of essential oils of *Anethum graveolens* L. and *Melaleuca cajuputi* Powell against *Rhyzopertha dominica* (F.) (Coleoptera: Bostrichidae). *International Journal of Tropical Insect Science*. <https://doi.org/10.1007/s42690-020-00359-6>
- Karataş T, Korkmaz F, Karataş A and Yildirim S. (2020). Effects of Rosemary (*Rosmarinus officinalis*) extract on growth, blood biochemistry, immunity, antioxidant, digestive enzymes and liver histopathology of rainbow trout, *Oncorhynchus mykiss*. *Aquaculture Research* 26:1533–1541. <http://doi.org/10.1111/anu.13100>
- Kumari S, Pundhir S, Priya P, Jeena G, Punetha A, Chawla K, Jafaree Z F, Mondal S and Yadav G. (2014). EssOilDB: a database of essential oils reflecting terpene composition and variability in the plant kingdom. *Database* 2014: 1–12. <http://doi.org/10.1093/database/bau120>
- Liu G L, Liu L, Hu Y and Wang G X. (2021). Evaluation of the antiparasitic activity of coumarin derivatives against *Dactylogyrus intermedius* in goldfish (*Carassius auratus*). *Aquaculture* 533: 736069. <http://doi.org/10.1016/j.aquaculture.2020.736069>
- Menses J O. *et al.* (2020). Acute toxicity of hot aqueous extract from leaves of the *Terminalia catappa* in juvenile fish *Colossoma macropomum*. *Aquaculture International* 28: 2379–2396. <https://doi.org/10.1007/s10499-020-00596-z>
- Meneses J O, do Couto M V S, Sousa N C, Cunha F S, Abe H A, Ramos F M, Chagas E C, Chaves F C M, Martins M L, Maria A N, Carneiro P C F and Fujimoto R Y. (2018). Efficacy of *Ocimum gratissimum* essential oil against the monogenean *Cichlidogyrus tilapiae* gill

- parasite of Nile tilapia. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia* 70(2): 497–504. <https://doi.org/10.1590/1678-4162-9667>
- Moghrovyan A, Sahakyan N, Babayan A, Chichoyan N, Petrosyan M and Trchounian A. (2019). Essential oil and ethanol extract of oregano (*Origanum vulgare* L.) from armenian flora as a natural source of terpenes, flavonoids and other phytochemicals with antiradical, antioxidant, metal chelating, tyrosinase inhibitory and antibacterial activity. *Current Pharmaceutical Design* 25: 1809–1816. <https://doi.org/10.2174/1381612825666190702095612>
- Mohammadi G, Rashidian G, Hoseinifar S H, Naserabad S S and Doan H V. (2021). Ginger (*Zingiber officinale*) extract affects growth performance, body composition, haematology, serum and mucosal immune parameters in common carp (*Cyprinus carpio*). *Fish and Shellfish Immunology* 99: 267–273. <https://doi.org/10.1016/j.fsi.2020.01.032>
- Nafiqoh N, Sukenda S, Zairin M, Alimuddin A, Lusiastuti A, Sarter S, Caruso D and Avarre J C. (2020). Antimicrobial properties against *Aeromonas hydrophila* and immunostimulant effect on *Clarias gariepinus* of *Piper betle*, *Psidium guajava*, and *Tithonia diversifolia* plants. *Aquaculture International* 28: 1–13. <http://doi.org/10.1007/s10499-019-00439-6>
- Nagi A A, Abdullah M I, Jasmani S, Zakariah M I, Karim N U and Hassan M. (2016). Acute toxicity and behavioural responses of African catfish, *Clarias gariepinus* to a methanolic extracts of *Melaleuca cajuputi* leaves. *Research Journal of Fisheries and Hydrobiology* 11(7): 1–6.
- Nero V, Farwell A, Lister A, Van Der Kraak G, Lee L E J, Van Meer T and Dixon D G. (2006). Gill and liver histopathological changes in yellow perch (*Perca flavescens*) and goldfish (*Carassius auratus*) exposed to oil sands process-affected water. *Ecotoxicology and Environmental Safety* 63(3): 365–377. <http://doi.org/10.1016/j.ecoenv.2005.04.014>
- Noor A A M, Yusuf S M, Abdul Wahab W N A W, Adam M F I C and Sul'ain M D. (2020). Evaluation on composition, antioxidant and toxicity of *Melaleuca cajuputi* leaves. *Advances in Traditional Medicine* 21: 693–699. <https://doi.org/10.1007/s13596-020-00479-x>
- Nunes, B., Antunes, S.C., Gomes, R., Campos, J.C., Braga, M.R., Ramos, A.S., Correia, A.T., 2015. Acute effects of tetracycline exposure in the freshwater fish *Gambusia holbrooki*: antioxidant effects, neurotoxicity and histological alterations. *Archives of Environmental Contamination and Toxicology* 68: 371–381. <https://link.springer.com/article/10.1007/s00244-014-0101-z>

- Pal S, Kokushi E, Koyama J, Uno S and Ghosh A R. (2012). Histopathological alterations in gill, liver and kidney of common carp exposed to chlorpyrifos. *Journal of Environmental Science and Health, Part B. Pesticides, Food Contaminants, and Agricultural Wastes* 47(3): 180–195. <https://doi.org/10.1080/03601234.2012.632285>
- Prado-Audelo M L D, Cortés H, Caballero-Florán I H, González-Torres M, Escutia-Guadarrama L, Bernal-Chávez S A, Giraldo-Gomez D M, Magaña J J and Leyva-Gómez G. (2021). Therapeutic applications of terpenes on inflammatory diseases. *Frontiers in Pharmacology* 12: 704197. <http://doi.org/10.3389/fphar.2021.704197>
- Rashidian G, Boldaji J T, Rainis S, Prokić M D and Faggio C. (2021). Oregano (*Origanum vulgare*) Extract Enhances Zebrafish (*Danio rerio*) Growth Performance, Serum and Mucus Innate Immune Responses and Resistance against *Aeromonas hydrophila* Challenge. *Animals*, 11(2): 299. <https://doi.org/10.3390/ani11020299>
- Rodrigues S, Antunes S C, Nunes Band Correia A T. (2019). Histopathological effects in gills and liver of *Sparus aurata* following acute and chronic exposures to erythromycin and oxytetracycline. *Environmental Science and Pollution Research* 26: 15481–15495. <https://doi.org/10.1007/s11356-019-04954-0>
- Rodrigues S, Antunes S C, Nunes Band Correia A T. (2017). Histological alterations in gills and liver of rainbow trout (*Oncorhynchus mykiss*) after exposure to the antibiotic oxytetracycline. *Environmental Toxicology and Pharmacology* 53: 164–176. <http://dx.doi.org/10.1016/j.etap.2017.05.012>
- Roszaini K, Nor Azah M A, Mailina J, Zaini S and Faridz Z M. (2013). Toxicity and antitermite activity of the essential oils from *Cinnamomum camphora*, *Cymbopogon nardus*, *Melaleuca cajuputi* and *Dipterocarpus* sp. against *Coptotermes curvignathus* K. *Wood Science Technology* 47: 1273–1284. <http://doi.org/10.1007/s00226-013-0576-1>
- Sahimi M B M K, Nagi A M, Hamdan N A, Zakariah M I, Yusoff N A H, Tosin O V and Hassan M. (2022). Cajeput *Melaleuca cajuputi* extract supplementation in diets of *Macrobrachium rosenbergii*: insight on the growth, immunological responses and resistance against *Aeromonas hydrophila*. *Aquaculture Research* 00: 1–12. <http://doi.org/10.1111/are.15851>
- Salome-Abarcaa L F, Mandronea M, Sannac C, Poli F, Hondel C, Klinkhamere P G L, Choi Y H. (2020). Metabolic variation in *Cistus monspeliensis* L. ecotypes correlated to their plant-fungal interactions. *Phytochemistry* 176: 112402. <https://doi.org/10.1016/j.phytochem.2020.112402>

- Septiana S, Yuliana N D, Bachtiar B M, Putri S P, Fukusaki E, Laviña W A and Wijaya C H. (2020). Metabolomics approach for determining potential metabolites correlated with sensory attributes of *Melaleuca cajuputi* essential oil, a promising flavor ingredient. *Journal of Bioscience and Bioengineering* 129(5): 581–587. <https://doi.org/10.1016/j.jbiosc.2019.12.005>
- Shivappa R B. (2015). Laboratory evaluation of safety and efficacy for Melafix (*Melaleuca cajuputi* extract). *Journal of Exotic Pet Medicine* 24: 188–192. <http://dx.doi.org/10.1053/j.jepm.2015.04.020>
- Strzyzewska E, Szarek J and Babinska I. (2016). Morphologic evaluation of the gills as a tool in the diagnostics of pathological conditions in fish and pollution in the aquatic environment: a review. *Veterinarni Medicina* 61: 123–132. <https://doi.org/10.17221/8763-VETMED>
- Tacon A G J. (2020). Trends in global aquaculture and aquafeed production: 2000-2017. *Reviews in Fish Science and Aquaculture* 28: 43–56. <http://doi.org/10.1080/23308249.2019.1649634>
- Ukit U, Widiana A, Rahmawati E, Hasby R M. (2019). Antibacterial activities test of Cajuput leaf waste extract (*Melaleuca cajuputi* Powell) on pathogenic bacteria. *Journal of Physics: Conference Series* 1402(3): 033030. <http://doi.org/10.1088/1742-6596/1402/3/033030>
- Valenzuela-Gutiérrez R, Lago-Lestón A, Vargas-Albores F, Cicala F and Martínez-Porchas M. (2021). Exploring the garlic (*Allium sativum*) properties for fish aquaculture. *Fish Physiology and Biochemistry* 47: 1179–1198. <http://doi.org/10.1007/s10695-021-00952-7>
- Verma R S, Mehta A and Srivastava N. (2007). *In vivo* chlorpyrifos induced oxidative stress: attenuation by antioxidant vitamins. *Pesticides Biochemistry and Physiology* 88(2): 191–196.
- Yancheva V S, Stoyanova S G, Valcheva I G and Georgieva E S. (2016). Histological response of fish gills to metal pollution: common carp, *Cyprinus carpio* L., and common rudd, *Scardinius erythrophthalmus* L., from Topolnitsa Reservoir, Bulgaria. *Acta Zoologica Bulgarica* 68(1): 103–109.
- Zhang X P, Li W X, Ai T S, Zou H, Wu S G and Wang G T. (2014). The efficacy of four common anthelmintic drugs and traditional Chinese medicinal plant extracts to control *Dactylogyrus vastator* (Monogenea). *Aquaculture* 420: 302–307. <http://doi.org/10.1016/j.aquaculture.2013.09.022>